

thority of Dr. Coues, who obtained one specimen.

71. *NETION CAROLINENSIS*. *Green Winged Teal*.—Dr Coues obtained one single specimen at Rigoulette.

72. *AIX SPONSA*. *Wood Duck*.—Not rare in interior. Breeds in hollow trees

73. *AYTHYA AMERICANA*. *Redhead*.—I saw a single specimen in the water at Baie des Roches, 23 September. Am told that it is common.

74. *CLANGULA ISLANDICA*. *Barrow's Golden Eye*.—Common in rivers as far as Natashquan. Said to occur in Esquimaux River in mild winters.

Nests of this species have been found in the woods near Lake Champlain, and the ducklings were seen commonly on the Godbout River during summer.—C.

75. *CLANGULA ALBEOLA*. *Buffle-head Duck*.—Common in fall.

76. *HAREIDA GLACIALIS*. *Long-tailed Duck*.—Common in mouths of rivers in spring and fall.

77. *HISTRIONICUS MINUTUS*. *Harlequin Duck*.—Rather rare. Mouths of rivers, spring and fall. Probably breeds

78. *SOMATERIA MOLLISSIMA DRESSERI*. *American Eider Duck*.—Abundant everywhere. Breeds.

79. *SOMATERIA SPECTABILIS*. *King Eider*.—Abundant in spring in large flocks. I shot a great many of them. It is said to breed in this region occasionally. In *The Canadian Sportsman and Naturalist*, vol. 1, No. 7, July 15th, 1881, p. 51, in an article headed "Bird-nesting in Labrador," Mr. Napoleon A. Comenau, the writer, whom I know personally and who spoke with me personally to the same purport, says that on a small island opposite Mingan: "Indeed, one small island, visited by us, was almost covered with the nests of this species (*S. mollissima*), and here we first found the nest of its congener, the King Eider (*S. spectabilis*)." "This is, I believe, the first record of this rare nest found on the Atlantic.

"Bird-nesting in Labrador" was written by Mr. William Couper, the Editor of this Journal.

*SOMATERIA V-NIGRA*. *Pacific Eider*.—Abundant in large flocks in spring. I myself obtained specimens that had the decided "V-shaped black mark" on the chin, and was told by the natives that there were "three different species of spring ducks so near alike that you could hardly tell the difference." This species has been doubted by several authorities. I still believe that I can secure specimens and prove its occurrence unquestionably.

80. *CEDEMIA AMERICANA*. *Black Scoter*.—Abundant. Breeds by inland ponds.

81. *MELANETTA VRIIVETINA*. *White-winged Coot*.—Common in fall, rare in spring. Not known to breed.

82. *PELIONETTA PERSPICILLATA*. *Sea Coot*.—Common in spring, rare in late fall. Not known to breed

83. *MERCUS MERGANSER AMERICANUS*. *Fish Duck*.—I have seen one specimen taken near Fort Island.

84. *MERCUS SERRATOR*. *Red-breasted Merganser*.—Common in spring and fall. Breeds occasionally.

85. *LOPHODYTES CUCULLATUS*. *Hooded Merganser*.—Rather rare but occasional.

86. *SULA BARRHANA*. *Garnet*.—Common in Gulf of Saint Lawrence. Occasionally seen near the Labrador coast.

87. *PHALACROCORAX CARBO*. *Common Cormorant*.—Abundant off Meccattin Islands. Breeds.

88. *PHALACROCORAX DIOPHUS*. *Double-crested Cormorant*.—Common with the former.

89. *STERCORARIUS POMATORHINUS*. *Pomarine Jaeger*.—I have seen a specimen of this species I think taken near the mouth of Esquimaux River. Dr. Coues also obtained it.

90. *STERCORARIUS PARASITICUS*. *Richardson's Jaeger*.—Shot a specimen in St. Lawrence River, about opposite Point des Monts.

91. *STERCORARIUS BUFFONI*. *Buffon's Jaeger*.—Seen by Dr. Coues.

92. *LARUS GLAUCUS*. *Burgomaster*.—Not rare. I obtained several specimens. Breeds.

93. *LARUS MARINUS*. *Great Black-backed Gull*.—Abundant and breeds all along the Labrador coast.

94. *LARUS ARGENTATUS SMITHSONIANUS*. *Herring Gull*.—Common. Breeds everywhere.

95. *RISSA TRIDACTYLA*. *Kittiwake Gull*.—Common in spring and fall. Breeds occasionally.

96. *LARUS PHILADELPHICUS*. *Nonparle's Gull*.—Common in large flocks in fall, perhaps spring, but not known to breed on the Labrador coast.

97. *STERNA MACHURA*. *Arctic Tern*.—An abundant spring and fall migrant in the Gulf. (I found nests of this Tern abundant on islands near Natashquan.—C.)

98. *STERNA FLUVIATILIS*. *Common Tern*.—Seen at Rigoulette by Dr. Coues.

99. *FULMAREX GLACIALIS*. *Fulmar*.—Recorded by Dr. Coues off Belle Isle.

100. *CYMORHENA LEUCORHONA*. *Leach's Petrel*.—Common off coast as far at least as to Belle Isle.

101. *PUFFINUS MAJOR*. *Greater Shearwater*.—Not rare off shore along the whole coast.