

To this notion of the existence of two groups of crystalline rocks similar in lithological character but different in age, we have to object that the hypozoic gneiss is identical with the Green Mountain gneiss, not only in lithological character, but in the presence of certain rare metals, such as chrome, titanium, and nickel which characterise its magnesian rocks; all of these we have shown to be present in the unaltered sediments of the Quebec group, with which Sir William Logan has identified the gneiss formation in question. Besides which the lithological and chemical characters of the Appalachian gneiss are so totally distinct from the crystalline strata of the Laurentian system, with which Prof. Rogers would seem to identify them, that no one who has studied the two can for a moment confound them. Prof. Rogers is therefore obliged to assume a new series of crystalline rocks, distinct from both the Laurentian and Huronian systems, but indistinguishable from the altered palæozoic series, or else to admit that the whole of his gneissic series in Pennsylvania is, like the corresponding rocks in Canada, of palæozoic age.\* We believe that nature never repeats herself without a difference, and that certain variations in the chemical and mineralogical constitution of sediments mark successive epochs so clearly that it would be impossible to suppose the formation in adjacent regions of a series of crystalline schists like those of the Alleghanies contemporaneous with the sediments which produced the Laurentian system. We have elsewhere indicated the general principles upon which is based this notion of

---

\* Dr. Bigsby in 1824 described an extensive tract of gneissoid rocks on Rainy Lake and Lake Lacroix, north of Lake Superior. The general course of the strata he states to be "from N. W. to N. by W., with a corresponding easterly dip;" but he elsewhere speaks of the gneiss as running (dipping?) E. N. E. This gneiss often contains beds and disseminated grains of hornblende, and passes in some places into micaceous, chloritic and greenstone slates, and syenite. Staurolite is abundant in the mica schists, and octahedral iron occurs in the chloritic slates. A porphyritic granite containing beryl is also met with in this region. This gneiss is regarded by Dr. Bigsby as belonging "to transition rocks, from its constant proximity to red sandstone, the oldest organic limestone, and trap." (*Am. Jour. Sci.*, (1) viii, 61). The lithological and mineral characters of these crystalline strata seem to be distinct from those of the Laurentian system, and to resemble those of the Appalachians. Too much praise cannot be ascribed to Dr. Bigsby for his early and extensive observations on the geognosy and mineralogy of British North America.