

inner spur of the anterior tibia stout, broad, bifurcate at apex; the outer spurs of the anterior tibia and those of the middle and posterior tibiae long, slender and simple: antennae stout, thicker at apex, the third segment one-third longer than fourth; clypeus deeply emarginate; wings smoky, darker at base; stigma and costa black; the marginal cross-vein originating in the lower posterior angle of the stigma, and joining the third submarginal cell at the end of the second third; posterior wings with two middle cells; the lanceolate cell much shorter than the submedian cell. Length, 10 mm.

Habitat, Olympia, Washington. 4 ♀♀, May 28, July 2. Trevor Kincaid, collector.

BIBLIOGRAPHY.

PERINEURA, Hartg.

1837. Hartig, Die Fam. Blattwes. und Holzwes., 303.

1867. Norton, Trans. Amer. Ent. Soc., I., 38.*

1868. Norton, Trans. Amer. Ent. Soc., II., 224.

1884. Konow, Deut. Ent. Zeit., 327.

1890. Konow, Deut. Ent. Zeit., 236.

Synairema, Hartig.

1837. Hartig, Die Fam. Blattwes. und Holzwes., 314.

1882. — *Perineura*, Hartg., Kirby, List Hymen. Brit. Mus., I., 279.

1886. Provancher, Faun. Ent. Can. Suppl., II., 15.

1887. Cresson, Trans. Amer. Ent. Soc., Suppl., 16, 19.

1890. — *Perineura*, Hartg., Konow, Deut. Ent. Zeit., 252.

Americana, Prov.

1885. *Synairema*, Provancher, Can. Ent., XVII., 50.

1886. *Synairema*, Provancher, Faun. Ent. Can. Suppl., II., 15.

1887. *Synairema*, Cresson, Trans. Amer. Ent. Soc. Suppl., 169.

Habitat, Canada.

Kincaidia, sp. nov.

Habitat, Olympia, Washington.

Pacifica, Prov.

1886. *Synairema*, Provancher, Faun. Ent. Can. Suppl., II., 15.

1887. *Synairema*, Cresson, Trans. Amer. Ent. Soc. Suppl., 169.

Habitat, Vancouver Island.

*The lanceolate cell is said to have an oblique cross-nervure, and *Synairema* to have this cell closed in the middle.