

widening before tip, tip black; antennæ 7-jointed, reaching to base of honey-tubes, annulated with brownish-red at joints; honey-tubes long, reaching beyond tip of abdomen, pale greenish, very slightly infuscated at tip; style short, conical, pale green; legs uniform pale green, feet very slightly infuscated.

Winged individual.—Length .06 inch. Elongate ovate and shining green. Vertex of head reddish; eyes brown; antennæ reaching beyond tip of abdomen, dark brownish black, excepting basal joints, which are pale; thorax shaded with brown; abdomen pale; legs pale greenish yellow, knees black; wings hyaline, veins greenish yellow, discoidal vein black.

4. *S. avenæ*, Fab.

5. "*viticola*, Thomas. First detected here early in March on wild grape vines growing in our swamps; later it becomes quite common on cultivated vines.

6. *S. setariæ*, Thos.

7. "*euphorbiæ*, Thos.

8. "*euphorbicola*, Thos.

9. "*asclepiadis*, Fitch. Very common here in early spring on *Asclepias cornuti*.

10. *S. erigeronensis*, Thos.

11. "*cereopsis*, Thos.

12. "*lactucæ*, Linn. Occasionally found here on lettuce.

13. "*polygoni*, Walker.

14. "*salicicola*, Thos.

15. "*verbenæ*, Thos.

16. "*rubi*, Kalt. Rarely found in early spring on under surface of leaves of *R. villosus*.

17. *S. pisi*, Kalt.

18. "*gerardiæ*, Thos.

19. "*heucherae*, Thos.

20. "*cucurbitæ*, Thos.

21. "*tanacetii*, Linn.

22. "*fragariæ*, Koch. Var. *immaculata*, Riley.

23. "*menthae*, Buckton.

24. "*absinthii*, Linn.

25. "*achyrantes*, Monell.

26. "*calendulla*, Monell.