

level as would have occurred without climate change; however, the cost of achieving this is unclear. Nonetheless, climate change may intensify difficulties in coping with rapid population growth. An increase or change in UV-B radiation at ground level resulting from the depletion of stratospheric ozone will have a negative impact on crops and livestock.

The rotation period of forests is long and current forests will mature and decline during a climate in which they are increasingly more poorly adapted. Actual impacts depend on the physiological adaptability of trees and the host-parasite relationship. Large losses from both factors in the form of forest declines can occur. Losses from wildfire will be increasingly extensive. The climate zones which control species distribution will move poleward and to higher elevations. Managed forests require large inputs in terms of choice of seedlot and spacing, thinning and protection. They provide a variety of products from fuel to food. The degree of dependency on products varies among countries, as does the ability to cope with and to withstand loss. The most sensitive areas will be where species are close to their biological limits in terms of temperature and moisture. This is likely to be, for example, in semi-arid areas. Social stresses can be expected to increase and consequent anthropogenic damage to forests may occur. These increased and non-sustainable uses will place more pressure on forest investments, forest conservation and sound forest management.

Natural terrestrial ecosystems

Natural terrestrial ecosystems could face significant consequences as a result of the global increases in the atmospheric concentrations of greenhouse gases and

the associated climatic changes. Projected changes in temperature and precipitation suggest that climatic zones could shift several hundred kilometres towards the poles over the next fifty years. Flora and fauna would lag behind these climatic shifts, surviving in their present location and, therefore, could find themselves in a different climatic regime. These regimes may be more or less hospitable and, therefore, could increase productivity for some species and decrease that of others. Ecosystems are not expected to move as a single unit, but would have a new structure as a consequence of alterations in distribution and abundance of species.

The rate of projected climate changes is the major factor determining the type and degree of climatic impacts on natural terrestrial ecosystems. These rates are likely to be faster than the ability of some species to respond and responses may be sudden or gradual.

Some species could be lost owing to increased stress leading to a reduction in global biological diversity. Increased incidence of disturbances such as pest outbreaks and fire are likely to occur in some areas and these could enhance projected ecosystem changes.

Consequences of CO₂ enrichment and climate change for natural terrestrial ecosystems could be modified by other environmental factors, both natural and man-induced (eg by air pollution).

Most at risk are those communities in which the options for adaptability are limited (eg montane, alpine, polar, island and coastal communities, remnant vegetation, and heritage sites and reserves) and those communities where climatic changes add to existing stresses.