

the dorsum fuscous, apex of the segments black, with an interrupted yellow basal lunule; apical segments entirely blackish-fuscous; wings hyaline, the costa yellow, pterostigma short, the sides a little oblique, black. Length 53–41 millim.; alar expanse 49–41 millim.; pterostigma 1 $\frac{1}{4}$  millim.

*Hab.*: U. S. Chicago, Minnesota, Indiana, Pennsylvania, New York, Mass.

**CALOPTERYX, Leach.**

**C. VIRGINICA**, Drury ed. Westwood.—**C. DIMIDIATA**, Rambur.

Brassy-green, shining; labium, antennae, thoracic sutures, pectus, venter, and feet black; abdomen with the dorsal stripe, venter with segments 8–10 yellow; wings narrow, hyaline, the base somewhat flavescent, the apex blackish-fuscous; pterostigma in the female snow-white. Length 50–52 millim.; alar expanse 70–72 millim.; pterostigma 2 millim.

*Hab.*: Hudson's Bay, U. S. Mass. and Southward.

**C. MACULATA**, Beauv.—**C. HOLOSERICCUS**, Burm.—

**C. PAPILIONACEA**, Ramb.

Brassy-green or blue, shining; labium, antennae, thoracic sutures, pectus, venter, and feet black; abdomen with a dorsal yellow stripe, 8–10 segments in the female; wings very broad, densely reticulated, black, sometimes with hyaline spots in the male, or clouded with fuscous, and fuscous at the apex, the female with a snow-white pterostigma. Length 38–48 millim.; alar expanse 63–65 millim.; pterostigma 2–2 $\frac{1}{2}$  millim.

*Hab.*: U. S. Chicago, Mass. and Southward. A species common and widely diffused.

**C. APICALIS**, Burm.

Brassy-green, shining; labium, antennae, thoracic sutures, and pectus black, or in the female in part flavescent; feet long, black, with long cilia; wings narrow, hyaline, or with the apex narrowly fuscous, pterostigma absent. Length 42 millim.; alar expanse 62 millim.

*Hab.*: U. S. Pennsylvania, Mass.