sensitiveness of the skin became more acute, and in about two months she looked like a different person, active, willing to walk, and without fatigue; appetite much improved; the entire tone of the nervous system was such as not experienced for years previous, and returned to her home in British Columbia, with every hope of many years of increased usefulness.

CASE III. Mr. B., æt. 28 years, weight 128 lbs., temperate. Has found of late that he tires readily, and awakes in the morning not sufficiently refreshed. No organic disease. For fully three years, experienced indications of debility, without any special cause except irregularity as to intestinal absorption, abdomen at times considerably distended with gas. Marked palor of face and lips, and an exsanguine condition of the body generally. A peculiar leaden feeling about the abdomen, with a ballooned duodenum. Bowels not regular. No nausea or vomiting. Appetite as a whole good, but the strength and support gained, not in proportion to food taken, evidencing the escape of nutrient material. Urine voided in normal quantity, but overcharged with lithates. About twelve months ago became indisposed, owing to the peculiar feeling in his bowels, which continued until May, 1898, when I placed him under neurotone treatment twice a week for two months, after which the improvement as to his general condition was most marked. Prior to that date, tonics had little effect. At present he can wheel 15 to 20 miles a day, without inconvenience, which he could not undertake for months previous. The tone of the entire nervous system is most marked, and life now most enjoyable.

CASE IV. Mrs. F., æt. 78 years, mother of a large family, weight 108 Temperate. No evidence of organic disease. Muscles thin and flabby. Functions of the system, regular as a whole. For several years experienced flatulence and constipation, regulated by occasional castor oil. About two years ago, the digestive system, particularly the alimentary canal, was the seat of very considerable irregularity. A degree of general debility followed, attended by marked weakness, almost approaching heart failure, and inability to move about, with accustomed activity. In May, 1898, placed her under neurotone treatment, continued twice each week, for a period of two months. At the expiration of that time, markedly improved, as to alimentary digestive power, and the general vigor of her system. Oct. 12, 1898. Moves about her home with the activity of twenty years ago, and now expresses herself as feeling almost youthful once more. This is an illustration of neurons, almost on the shelf, called into action again and becoming useful factors in the promotion of renewed life and activity.