

- Largidea opaca* Uhl. (MS.) Smaller and slighter than New York examples.
- Campptobrochis grandis* Uhl. Common. Apparently of nocturnal habits, as I took several flying around the candle at our camp; some of these individuals were extremely dark, even approaching an almost uniform piceous black; others were as pale as those taken near Buffalo.
- Ncoborus Petitii* Uhl. (MS.) Several pale examples.
- Fulvius anthocoroides* Uhl. One example.
- Monalocoris filicis* Linn. Common everywhere on ferns.
- Hyaliodes vitripennis* Say. This neat little species was taken frequently on pines, and occasionally on other trees and bushes.
- Pilophorus amœmus* Uhl. Common on pine trees.
- Globiceps flavomaculatus* Fab. One example taken. This species was erroneously cited as occurring at Buffalo, in my list of Capsidæ from that locality (CAN. ENT., xix., p. 72, 1887). The insect there referred to was the next, which superficially resembles the *flavomaculatus*.
- Mimoceps gracilis* Uhl. (MS.) A few swept from grass and weeds near a rivulet among the hills.
- Mecomma gilvipes* Uhl. (MS.) Three examples taken with the preceding.
- Macrolophus seperatus* Uhl. Three examples.
- Stiphrosoma stygica* Say. Scattering.
- Labops hesperia* Uhl. A few brachypterous examples swept from an oat field on the flats near the mouth of the river.
- Idolocoris agilis* Uhl. Common on the lowlands.
- Orthotylus alternatus* Uhl. Common on bushes, especially the hazelnut.
- Macrotylus guttatus* Uhl. (MS.) One example of this elegant little Capsid was swept from briars growing on a rocky hillside near the river.
- Rhinocapsus Vanduzeei* Uhl. (MS.) Not uncommon among rank weeds in damp situations.
- Psallus variabilis* Fall. A single example.
- Psallus antennatus* Uhl. (MS.) Several taken.
- Plagiognathus obscurus* Uhl.