

4. July.

m pubes-
mentosis,
lycis laci-
oribus.—

the sea-
ne—Aug.
ut petals;
going spe-

canescens;
cano-to-
nfertiflo-
lyce triplo

ly. v. s.

lternis ob-
ibus utrin-
cis calyci-
s acutis pe-

June—
han in any

t. terol. 15.

1778. t. 2.
½ May,
icate little
arge tracts
s common

427. TALINUM. *Juss. gen. pl. 346.*

1. *T. foliis cylindricis carnosis, corymbis terminalibus pedunculatis.* — *teretifolium.*
On sunny rocks: Delaware and Virginia. 4. July.
v. v. Flowers purple.

428. PORTULACCA. *Gen. pl. 824.*

1. *P. foliis cuneiformibus, floribus sessilibus.* — *Willd. sp. oleracea.*
pl. 2. p. 859.
Icon. *Schkuhr handb. 130.*
Purslane is a common weed in gardens.

429. CHELIDONIUM. *Gen. pl. 880.*

1. *C. foliis alternis pinnatis lobatis, umbellis axillaribus pedunculatis.* — *Willd. sp. pl. 2. p. 1141.*
Icon. *Fl. dan. 542. Engl. bot. 1581.*
Near old buildings and among rocks: New England and Pennsylvania; probably introduced. 4. May, June. *v. v.* The yellow fluid which it affords by breaking the stems, is considered an infallible remedy in curing warts and corns.
2. *C. caulis in summitate geminatim diphylloides, foliis sessilibus lobato-pinnatifidis, pedunculo solitario unifloro, capsula oblongo-ovata.* — *Mich. fl. amer. 1. p. 309.*
In shady woods, near rivulets: Kentucky and Tennessee. *Michaux.* 4. May. *v. s.* Resembles *Sanguinaria* very much, and seems to be the connecting link to both genera.

430. GLAUCIUM. *Juss. gen. pl. 261.*

1. *G. caule glabro, foliis caulinis amplexicaulibus repandis, pedunculis unifloris, siliqua tuberculato-scabriuscula.*
Smith. fl. brit. 2. p. 563.
Chelidonium Glaucium. — *Willd. sp. pl. 2. p. 1142.*
Icon. *Fl. dan. 585. Engl. bot. 8.*
On the sea-coast of Virginia and Carolina. 6. June, July. *v. v.* Flowers large, yellow.

431. PAPAVER. *Gen. pl. 681.*

1. *P. capsulis hispidis, scapo unifloro nudo hispido, foliis subpinnatis, foliolis lanceolatis; inferioribus subincisis.*
— *Willd. sp. pl. 2. p. 1145.*