

(4) If the fall has been very gradual, it may occasionally happen that the respiration stops completely, and still the blood pressure rises again, the respiration recommencing spontaneously in the course of the rise. In the same way, when the inhalation has been discontinued, the respiration may stop during the after-fall of the blood pressure and begin again spontaneously. As a rule, if the respiration has stopped, or even becomes slow and feeble at the time when the inhalation is discontinued, and artificial respiration is not resorted to, the fall in blood pressure will continue until death ensues.

(5) There are two conditions which frequently disturb the gradual fall of the blood pressure—viz., struggling and holding the breath,—and it is only by great care that they can be avoided in animals.

(6) Struggling, independently of any change in the respiratory rhythm, appears generally to raise the blood pressure. In one case of a dog much weakened from phosphorus the pressure fell every time he struggled.

(7) When struggling is accompanied, as it often is, by acceleration of the respiration and pulse, especially if the respiration is deep and gasping, it leads to a more rapid inhalation of chloroform, and consequently to a more rapid fall of blood pressure and a greater after-fall. In order to keep the chloroform cap or inhaler in its place during the animal's struggles, the administrator is obliged to hold it down more tightly over the nose and mouth, and this materially assists in hastening the rapidity of the inhalation, and consequently of the fall in blood pressure.

(8) The effect of involuntarily holding the breath—which, as anybody can prove by experiment upon himself, must happen when an inhaler saturated with chloroform is first applied to the face—is much more remarkable, the pressure often falling with great suddenness, while the heart's action is markedly slowed. As soon as the animal draws breath again, the pressure rises as suddenly as it fell, but the gasping respiration which succeeds then causes very rapid inhalation of chloroform, with immediate insensibility and a rapid fall of blood pressure, which quickly becomes dangerous.

(9) The combination of struggling with alternate holding the breath and gasping, which results if chloroform is applied closely to the face without sufficient dilution with air, causes violent fluctuations, and then a speedy fall of the blood pressure, which very soon leads to a dangerous depression with deep insensibility and early stoppage of the respiration. The after-fall under these circumstances is rapid and prolonged. It is this combination of events which causes struggling animals to go under chloroform so quickly.

(10) The effect of holding the breath may occasionally cause a temporary fall of blood pressure

after the chloroform inhalation has been stopped, or even when the animal is quite out of chloroform. This fall is recovered from directly the animal breathes again.

(11) Slight continuous asphyxia, such as is produced by pressure on the neck by straps, a badly-fitting muzzle, or hindrance of the chest movements by the legs being too tightly bound down, gives rise to exaggerated and irregular oscillations of the blood pressure, and slowing and irregularity of the heart's action. If it leads to, or is accompanied by deep gasping inspiration, it is apt, like anything else which causes this, to increase the intake of chloroform and bring about a rapid decline of blood pressure.

(12) Complete or almost complete asphyxia, as by forcibly closing the nose and mouth or closing the tracheal tube after tracheotomy, has an effect similar to, but more marked than, that produced by holding the breath, and the character of the trace corresponds precisely to that produced by irritation of the peripheral end of the cut vagus. The pressure falls extremely rapidly, sometimes almost to zero, and the heart's action becomes excessively slow, or even stops for a few seconds. If the Fick trace of Experiment 148 be compared with the photographic reproduction of Trace A of the Glasgow Committee, it will be seen that they are identical, and that the slow action of the heart with great fall of pressure, which the Glasgow Committee attributed to some capricious action of chloroform upon the heart, was undoubtedly due to asphyxia.

(13) This effect of asphyxia is the result of stimulation of the vagi. The proof of this is (a) that the trace corresponds exactly, as stated above, to that produced by direct irritation of the vagus, (b) division of both vagi entirely abolishes it, and (c) the administration of atropine which paralyses the vagus also abolishes it.

(14) In Trace 158 (Fick 4), which was taken during asphyxia after a full dose of atropine, it will be seen that there is an alternately slow and rapid pulse according to the phase of the respiratory movement, but no continued slowing of the heart as in vagus irritation. But there was still a distinct fall of pressure after the atropine when the breath was held, and it was thought that the slowing of the pulse above noted in this condition might be due to the disturbance of the heart from tension in the pulmonary vessels in the absence of respiratory movement, rather than to irritation of the vagi. To test this point Experiment 184 was instituted. In this experiment the dog's chest was forcibly inflated with bellows connected by a tube with the trachea, and the effect of this proceeding was to cause a fall of pressure and slowing of the heart exactly the same as involuntary holding of the breath. The dog was then poisoned with atropine, after which inflation of the chest