locality, Georgia. This is evidently different from any species I have seen.

Andrena nivalis, Smith, 1853.

\(\varphi\): "Tubercle \(\frac{\top}{\top}\); antennal joint 3 longer than 4+5; area rugose at base; abdomen distinctly punctured, but on an acciulated surface."

Type locality, Hudson's Bay. This is very near to \(A.\) semirufa, Ckll., from New Mexico; possibly it is the same species.

\(\frac{\top}{\top}\)

\(\text{New New Mexico}\); possibly it is the same species.

\(\text{Surface}\)

\(\text{Tubercle}\)

\(\text{Tube

Andrena hilaris, Smith, 1853.

? . "Tubercle of labrum wide, truncate, below its centre a projecting pencil of golden hairs; clypeus coarsely punctured, with a smooth shining carina down its middle; metathoracic area finely rugulose, with some coarse short longitudinal strige at its base; abdomen distinctly punctured, its surface also microscopically (hardly visibly) rugulose; wings very yellow." Type locality, Georgia. Evidently distinct from anything I have before me.

Andrena vicina, Smith, 1853.

Q. Length "about 14 mill.; tubercle wide, truncate; clypeus with wide unpunctured central elevation; area dull rugulose; abdomen finely rugulose all over, and distinctly but shallowly punctured." Smith gives the length as only 5 lines, and says the species closely resembles A. nitida. I hardly know what to make of this; the species referred to vicina in American collections has the tubercle low and rounded, by no means truncate; also, it does not have the very shiny, oval abdomen of nitida, which species, it may be remarked, has the tubercle concavely truncate. A. vicina is said to be from the United States and Nova Scotia.

Andrena placida, Smith, 1853.

9. "Tubercle very narrowly truncated; area rather coarsely rugulose, but not margined, I think; abdomen rugulose, a few sparse punctures; length about 12 mill., much smaller and narrower insect than vicina—looks like a parviceps." From "U. S." Smith gives the size the same as that of vicina, but this must be a mistake. A. placida seems to come close to A. Macgillivrayi.

Andrena fastuosa, Smith, 1879.

- ♀. "Tubercle / ✓ , difficult to say if actually emarginate; area and abdomen as ♂."
- †. "Tubercle emarginate; huge stipites! area rugose, unmargined;
 abdomen closely punctured." From Orizaba, Mexico. Andrena

 Andrena

 **Tubercle emarginate; huge stipites! area rugose, unmargined;
 abdomen closely punctured."

 **Tubercle emarginate; huge stipites! area rugose, unmargined;
 abdomen closely punctured."

 **Tubercle emarginate; huge stipites! area rugose, unmargined;
 abdomen closely punctured."

 **Tubercle emarginate; huge stipites! area rugose, unmargined;
 abdomen closely punctured."

 **Tubercle emarginate; huge stipites!
 **Tubercle emarginate