

- Pubescence above and nervures and stigma whitish *pruinus*.
6. Dark blue *cæruleus*.
Greenish ; abdomen usually more or less testaceous *zephyrus*.
7. Abdomen and tibiæ almost entirely testaceous *zephyrus*.
Abdomen not testaceous ; tibiæ black except often at base and
apex 8.
8. Sides of metathorax and pleura distinctly punctured, the latter
beneath with a distinct fovea ; mesonotum smooth, shining,
coarsely punctured *foveolatus*, sp. nov.
Sides of metathorax and pleura not distinctly punctured 9.
9. Vein III₅ and beyond almost obsolete ; head much longer than
wide ; mesonotum finely rugose, opaque, sparsely, finely punctured ;
antennæ short ; tarsi pale *coreopsis*, sp. nov.
Vein III₅ and beyond ordinary ; head not, or hardly, longer than
wide 10.
10. Mesonotum finely punctured 12.
Mesonotum coarsely punctured, shining 11.
11. Wing white, nervures and stigma white *albipennis*.
Wing ordinary, nervures and stigma dark ; metathorax coarsely
reticulated, with a semicircular enclosure bordered by a sharp
edge *Cressonii*.
12. Mesonotum shining ; head wider than thorax ; metathorax at apex
gibbous, smooth, shining ; nervures and stigma dark ; abdomen
subclavate, almost impunctate, usually darker towards apex ;
length 4 mm *sparsus*, sp. nov.
Mesonotum opaque ; abdomen hardly subclavate 13.
13. Abdomen bronze black, minutely punctured, bare impunctate apical
margins of segments broad ; nervures and stigma dark *obscurus*.
Abdomen less black, distinctly punctured, bare impunctate apical
margins of segments narrow, often pale testaceous ; nervures and
stigma pale : length, 4-6 mm *versatus*, sp. nov.

PARALICTUS, Rob.

Females.

- Cheek regularly rounded ; face narrowed below *simplex*.
Cheek with rounded angle below middle of eye ; face narrowed
below *platyparius*.
Cheek with rounded angle a little above middle of eye ; face not
narrowed below *cephalicus*.