

397. STRUTHIOPTERIS, Willd. - OSTRICH-FERN.  
920 *S. Germanica*, Willd. Common in rich alluvial soil, Kent Co., *Fowler*; very abundant on low grounds of Tobique, mouth of St. Francis, *Hay*, Bulletin II. 25; on Tobique River, *Dr. Bailey*, Can. Nat. 1864, p. 83.
398. ONOCLEA, L. - - - SENSITIVE-FERN.  
921 *O. sensibilis*, L. Rather common in northern counties, *Fowler*; common on St. John and tributaries, Campbellton, *Hay*, Herb.; along Tobique River, *Dr. Bailey*, Can. Nat. 1864, p. 83, 85.  
var. *obtusilobata*, Torr. Richibucto, *Fowler*; Havelock, King's Co., *Brittain*.
399. WOODSIA, R. Brown. - - - WOODSIA.  
922 *W. Ilvensis*, R. Br. Restigouche on Sugar Loaf and at mouth of Upsalquitch, *Fowler*; St. Stephen, *Vroom*; found at Keswick, Nashwaaksis and near Green Head, St. John, *Hay*; in Herb. *Mrs. Heustis*; Minister's Face, Kennebecasis, Bulletin II. 29; very abundant about Lower Tobique, *Hay*.  
923 *W. hyperborea*, R. Br. Rare. At tunnel in Restigouche, *Fowler*; Aroostook Falls, *Hay* and *Wetmore*, Bulletin II. 29, Herb.; Grand Falls, *Peter Jack*. This fern was given as *W. glabella* in former list through a mistake in copying the label.
400. DICKSONIA, L'Her. - - - DICKSONIA.  
924 *D. punctilobula*, Kunze. Very common in pastures, roadsides and open woods.
401. OSMUNDA, L. - - - FLOWERING-FERN.  
925 *O. regalis*, L. Swamps and wet places. Bass River, Fredericton, *Fowler*; Upper Tobique and Campbellton, *Hay*, Herb.; St. John, Herb. *Mrs. Heustis*; Tobique, *Dr. Bailey*, Can. Nat. 1864, p. 83.  
926 *O. Claytoniana*, L. Very common throughout. Lancaster, *Hay*, Herb.  
927 *O. cinnamomea*, L. Very common in swamps in northern counties, *Fowler*; Campbellton, *Hay*, Herb.  
var. *frondosa*, Gray. Beautiful specimens showing every step of the change from the common sterile form of frond to forms having several of the upper pinnæ completely fertile were once found by the writer at Molus River, Kent Co.