

finely punctured; antennæ about half as long as the body, rather slender, enlarging moderately toward tip; prothorax finely but strongly, closely punctured; elytra parallel, about two-thirds longer than wide, the punctures strong and rather sparse, the surface strongly shining, each broadly but rather abruptly swollen near the scutellum. Length, 1.4 mm.; width, 0.75 mm. New York, (near the city)

fusciceps, n. sp.

Darker brown, the head blackish; integuments much duller in lustre, the punctures denser, those of the elytra smaller, but more close-set; antennæ almost similar; prothorax relatively larger, strongly rounded at the sides, slightly narrower than the head; elytra rather more elongate and less subtruncate at apex, the swellings near the scutellum feebler and more diffuse. Length, 1.5 mm.; width, 0.7 mm. Wisconsin (Bayfield). Mr. Wickham.....*congener*, n. sp.

8. Eyes small, moderately prominent, the tempora behind them larger than in any other species of the genus, and from more than one-half to two-thirds as long as the eyes; front broadly convex, closely punctate in the male, sparsely in the female; male antennæ about half as long as the body, the five outer joints very faintly larger, those of the female two-fifths as long as the body, with the five outer joints more distinctly enlarged; prothorax densely punctate, wider than long, only slightly narrower than the head, narrowed anteriorly; elytra parallel, obtusely rounded at tip, finely, strongly and closely punctured, more elongate in the male and about three-fourths longer than wide; size small as in *piceus*. Wisconsin (Bayfield). Mr. Wickham. [= *Tanilotes lacustris*, Csy.].....*Wickhami*, Csy.
- Eyes larger, almost attaining the base; size much larger, about 2 mm. in length, similarly blackish in coloration and only moderately shining, strongly and closely punctured; antennæ barely two fifths as long as the body, the five-jointed club very broad and conspicuous; elytra parallel, nearly twice as long as wide, obtuse at apex; femoral tooth of the female much larger than in *Wickhami*. Pennsylvania (near Philadelphia).....*densus*, Csy.
9. Eyes separated by distinctly more than their own width; antennæ shorter, two-fifths as long as the body, gradually but strongly incrassate distally; elytral punctures coarser and less dense, the surface somewhat shining. Florida (Crescent City).....*Floridanus*, Csy.
- Eyes separated by not more than their own width and sometimes less; elytral punctures smaller, dense, the surface very dull; antennæ much longer, fully one-half as long as the body..... 10
10. Basal thoracic impressions distinct; form stouter, nearly as in *piceus*; antennæ thick, rather more than half as long as the body, only just