

Therapeutic Suggestions.

CYSTITIS.—There is an advantage, according to the *Jour. de Med. de Paris*, in prescribing salol internally in the form of an emulsion instead of powder form in the treatment of cystitis. The following formula is recommended :

R. Salol.....	gr. xxx-dr. i	2-4	
Pulv. tragacanthæ.....	gr. ii		12
Pulv. acaciæ.....	gr. xxx	2	
Tinc. simp.....	oz. ss	15	
Aq. distil. q. s. ad.....	oz. ii	60	

M. Sig.: Ft. emulsio. Sig.: One teaspoonful before each meal.
The following combinations are recommended by *Merck's Archives* in the treatment of cystitis :

To relieve the pain :

R. Ext. opii.....	gr. vi	40
Ext. hyoscyami.....	gr. v.	30
Olei theobrom. q. s.		

M. Ft. suppos. No. vi. Sig.: One to be introduced into the rectum at night; or :

R. Pulv. opii.....	gr. xii.	75
Camphoræ.....	dr. ss	2
Ext. belladonnae	gr. iii	20
Olei theobrom. q. s.		

M. Ft. suppos. No. vi. Sig.: One to be inserted at bedtime.
As an antiseptic and to relieve the irritation the following is of service :

R. Ext. belladonnae fl	gtt. xx	1 30
Sodii boratis.....	dr. ii	8
Acid benzoici.....	gr. xx	30
Tinc. opii camph.....	oz. iss	45
Olei gaultheriæ.....	gtt. xii	75
Syrupi.....	oz. ii	60
Aq. destil.....	oz. xii	120

M. Sig.: One dessertspoonful in water four times a day.

—*Los Angeles Medical Journal.*