-THE MANITOBA AND WEST CANADA



A J. urnal of Medicine, Surgery, Physiology, Chemistry, Materia Medica and Scientific News, being the journal of the Winnipeg and Manitoba Medical Associations. Published Monthly. Subscription 81 per annum in advance.

Vol. 6.

WINNIPEG, AUGUST, 1898.

No. 4.

FREE for a POSTAL

Desirous that every physician may have opportunity to make-trial of

DUNCAN FLOCKHART & CO'S

FLEXIBLE CAPSULES

I am instructed by Messrs. D. F. & Co. to send working samples to every physician making application for same. List of Capsules will be forwarded on request.

R. L. GIBSON, ^{86 Wellington St., West;} TORON'PO.

Lactopeptine Tablets

Same formula as Lactopeptine Powder. Issued in this form for convenience of patient—who can carry his medicine in his pocket, and so be enabled to take it at regularly prescribed periods without trouble.

"Everything that the science of pharmarcy c u do for improvement of the manufacture of Peps'n, Pancreatine, and Diastase, has been quietly applied to these ferments as compounded in Lactopeptine."

THE MEDICAL TIMEBAND HOSPITAL GABRITE.

Can be ordered through any Druggist. Samples free to medical men.

New York Pharmacal Association,

88 Wellington St., West, Toronto.

WAMPOLE'S ANTISPEPTIC SOLUTION

FORMOLID

16 oz Bottle 50c.

DIRECTIONS.

EXTERNALLY.

Dental. A non poisonous, palatable Antisertic and Germicide, useful in all stages of opertive work. Non-irritant. FORMOLID, pure or diluted, as indicated.

Surgical. For wounds, cuts, bruises, burns and scalds, etc. Prevents suppuration. Formolid, 1 part; water, 1 to 10 parts.

Mouth Wash. To preserve the teeth, sweeten the breath, harden the gums, etc. Formolid, 1 part; water, 4 to 20 parts.

Gargle, Nasal Application, etc. For inflamed mucous surfaces. Prophylactic. Formalid, 1 part; water 2 to 10 parts.

Vaginal Douche. Deodorant, Germieide. For Leucorrinea. Vaginitis, etc. Formolid, 1 part; water, 5 to 20 parts.

General. A healing antiseptic. Personal Hygiene. Excessive perspiration. Formolid, 1 part; water, 1 to 20 parts.

INTERNALEY.

In fermantative dyspepsia, diarrhœa, cholera Infantum, cholera morbis, etc.

Dose. One-half to two teaspoonfuls in water three or four times daily, as indicated. Can be given in combination with other medicines, except digestive terments.