

thorace sat longo concinne punctato, nucha tantum linea apicali elevata indicato, carina media elevata, plica sat distincta vix arcuata, interdum costula transversa instructo. Alæ fere vitreæ, parum pubescentes, haud longæ nec latæ, cellula basali et speculo magno nudis, stigmatibus tenui metacarpo sesqui longiore, radio sub angulo sat acuto egrediente, clava parva quam metacarpo vix brevior, disco macula magna fumata ad radii clavam usque extensa et ad marginem inferum fere descendente. Abdomen cupreo-aureum, splendens, segmentis 20 viridi, subtus interdum pallido et 30 40 que fere glabris, 3-6 sensim brevioribus, 80 parvo sub-erecto; ventre convexo sed haud compresso. Pedes validi rufescentes, coxis anterioribus interdum concoloribus. Mesosterni epimera punctata discreta, præsterna haud parva.

"Mas differt: antennis corporis fere longitudine articulis 20 brevior, 5-11 linearibus nigris longe hirtis, clava 2-articulata elongato-acuminata; metathorace longiore costula transversa magis distincta, abdomine oblongo macula pallida, femoribus basi et coxis totis viridiæneis.

"Sällsynt men troligen utbredd från Skane till Lappland."^a

(Thomson, 1878, pp. 158-159).

As Thomson, in the description of the species just given, makes *Pteromalus maculipennis* Walker a possible synonym, the original description of that species is appended to the foregoing, for the sake of convenience. It is impossible to decide this from the descriptions. *A. punctatus* was designated by Ashmead in 1894 as the type of the genus, and again as such, by the same author, in 1904. Further than in the places already cited, the species has not been mentioned again in the literature. The original description of *Pteromalus maculipennis* Walker follows:

"*Pteromalus maculipennis* Walker.

"Corpus mediocre; caput thorace paulo latius: antennæ clavatæ, crassæ, corporis dimidio longiores; articuli 50 ad 10 mm. longitudine decrescentes latitudine increscentes; clava articulo 100, paulo latior vix duplo longior; thorax ovatus; prothorax brevis: mesothoracis parapsidum suture vix conspicuæ: metathorax brevis: abdomen ovatum, subtus angulatum, apice elevatum acuminatum, non compressum, thorace paulo longius et latius: segmentum 1 mm. magnum: sequentia breviora; oviductus occultus; alæ mediocres: nervus cubitalis radialis vix brevior.

a. Rarely, but probably scattered (*utbredd*) from Servia to Lapland.