

The denuded wings were neatly mounted by Mr. Dimmock on white cards, to which they had been gummed. An interesting collection, illustrating the nerve structure of many of the genera of moths, was exhibited by him, to the great gratification of all present.

In the compilation of these memoranda in regard to the meetings of the Entomological Club, we are greatly indebted to the Secretary, Prof. C. V. Riley, who very kindly placed his notes at our disposal; also to Mr. B. P. Mann, of Cambridge, who did us similar service.

ON CATOCALA VERRILLIANA, WITH NOTES ON CATOCALA RELICTA.

BY A. R. GROTE, BUFFALO, N. Y.

Catocala Verrilliana Grote, Bul. B. S. N. S., 3, 12 (Aug., 1875).

The smallest N. Am. red-winged species known. It has a resemblance to *polygama* and *fratercula* in the ornamentation of the primaries. Fore wings gray, shaded with blackish; lines black. A diffuse basal black shade. T. a. line diffusely shaded with black. Reniform small, yellowish, more or less distinctly double ringed. Sub-reniform small, yellowish, disconnected with the t. p. line. T. p. line shaped much as in *polygama*. Hind wings bright red. Median black band narrow, tolerably even throughout, not attaining internal margin. Terminal band black, narrow, not quite attaining anal angle. A red apical mark, opposite which the fringes are pale. Medially the black band is five times outwardly scalloped, and opposite this scalloping the fringes are blackish. The red color again prevails below them on the margin, with the pale fringes; towards anal angle the fringes are dusky. Beneath, both wings red, with constricted median band on hind wings not attaining the margin.

Expanse 48 m. m. G. W. Belfrage, Bosque Co., Texas, June 13. Dedicated to Prof. A. E. Verrill, of New Haven. Interesting as a form of the red-winged group, resembling the yellow-winged and smaller species of the genus.

I am indebted to Prof. Hopffer, of the Royal Museum of Berlin, for a beautiful water colored drawing of the Texan *Catocala frederici* Grote,