

86. *Luscitiosa Clem.* N. Y., Maine, N. J.
87. *Lugens Walk.* Western States.
Eremitoides Streck.
88. *Eremitus Hübn.* Can., southward.
89. *Separatus Neum.* —
90. *Dollii Neum.* —
91. *Elsa Streck.* —

Genus *Dilophonota* Burm.

92. *Ello Linn.* West Indies, northward.
93. *Melancholica Grote.* West Indies, northward.
94. *Merianae Grote.* West Indies, Mex., Tex.
95. *Festa Hy. Edw.* —
96. *Obscura Fabr.* Southern States.
97. *Edwardsii Butler.* —

BROTIS VULNERARIA HUBN.

BY GEO. D. HULST.

In the CANADIAN ENTOMOLOGIST, this volume, p. 72, Mr. Ph. Fischer tells us of the capture of the above moth at Buffalo, N. Y., adds something of the bibliography of the species, and remarks upon its aberrant appearance and character. Permit me to add a few words to what was there said.

Guenee, Phalenites, vol. ii., p. 116, 1857, describes this species of Hubner, under the generic name *Sphacelodes*, crediting the species to Hubner, but ignoring his genus, as was Guenee's custom. He also describes the ♀, and pl. 22, f. 9, gives a figure of it. It differs very much from the ♂, having ciliated antennæ, and lacking the triangular flesh-colored costal spot. He is as much in doubt as was Hubner, as to its classification, but places it among his Fidoniidae. Walker, Cab. Brit. Mus. Geometridae, p. 213, 1860, catalogues *vulneraria* under *Brotis* Hubn., placing *Sphacelodes* Guen. as a synonym of the genus. He also is in doubt as to the proper location of the genus, but places it at the end of the Ennominae, and says it does not seem to fit well anywhere. Guenee's