

SYMPTOMS OF HOG CHOLERA— TREATMENT.

**You Must Follow These Special Directions
If Your Hogs Have the Hog Cholera.**

First symptoms are great debility and capricious appetites. You should separate the sick and apparently healthy at once; delay is very dangerous. Kill and burn those that are too sick to eat. It is extremely poor policy and dangerous to waste time on hogs that are too sick to eat.

Genuine malignant Hog Cholera cannot be fooled with, and demands heroic treatment without a moment's delay. Guard infested pens against all other animals. Do not go direct from the sick to the ones apparently healthy. Disinfect pens, nests, troughs, buckets, etc., with "International Pheno-Chloro," four tablespoons mixed with one bucket of water. A teaspoonful of "International Pheno-Chloro" mixed in every bucket of water or swill once per day will be very beneficial. Give them pure water. Feed the hogs that can eat easily digested food with two or three times the usual amount of "International Stock Food" mixed with slop and keep it up regularly four to five times per day for two or three weeks. Men of good judgment will not expect miraculous results, because they know that a diseased system cannot be changed nor a weak system thoroughly strengthened in three or four days. One 25-lb. pail of "International Stock Food" will not cure a herd of hogs nor show wonders if fed promiscuously to all your stock.

SUMMER FATTENING.

Summer and early fall is the ideal time to fatten hogs because they will make a more rapid gain. It takes a certain amount of food to supply the daily animal waste. The days of growth should be shortened so as to avoid the unnecessary waste. Market your hogs at six to eight months of age and you make a greater per cent of profit than your neighbor, who keeps his ten or twelve months, because you avoid 60 to 120 days of daily animal waste.