adian

me of ding.

-Sept

Oct.

Aug.

ept

ot .

- The their chief cha	racters—Co	ntinued.		
Colour, Size, Arrangement of Flowers and other Characters of the Plant,	Method of Propagation and Distribution	on of G	lacu rowth, roducts ured.	Methods of Eradication.
Pod 1 1				
Red, much bearded inside, 3-in dense terminal and axillary spikes berry reddish, 3-in.	stocks.	un-Newly broot-summer and pas		Break early, sum mer fallow.
White, small, in large terminal pani cles.			lds, past-	Summer fallow, cultivate.
Bright yellow; 3-in.; whole planglutinous; bud bearing large drop of white resin.	Seeds, wind,	in Fields, road sid	pastures, M	Iow, cultivate.
Yellow, head large, 1-sided	Seeds, runnir root-stock	g Grain field s, mer fallo	ds, sum-P	lough and culti-
Bright yellow; dense cormybs; leaves		Low land		6
White; ‡-in., crowded on spreading branches. White: heads very numerous are all		. Urain field	ures. s, sandy	
White; heads very numerous, small, crowded in a slender erect wand-like panicle.		grain fiel	us.	ummer fallow early, cultivate fall and spring.
White, tinged with purple, ½-in.; corymb; leaves coarsely toothed.	Seeds, wind, in hay.	Grain field dens.	ls, gar-Cı	litivate fall and
White; ½-in., panicled corymb; leaves entire or nearly so, rough.	"	. "		"
Rose pink; 3-in., showy, corymb; leaves clasping.	"	Low lands, pastures,	fields,	•
White, 1-in., a small crowded cluster; sflowering stern erect, naked.	and rutifiels.	Pastures	Bre	eak up sod, cul-
White, 1-in., many in a terminal S corymb; stem leafy, white, downy.	eeds, running root-stooks.	Meadows and	d past-	" ,
clusters; stem leafy; whole plant	wind. eeds, wind	**		11
fragrant. neonspicuous; terminal leafy clus-St	eeds, floods	Low ground,	fields. Dra	in thoroughla
aconspicuous Lin hanging at a				
leaves less than 1-in., rough, oblong linear, entire, opposite below, alternate above.	ing stems.	oram neice.	cu	mer fallow, ltivate con- antly.
	floods.	Grain crops,		
ellow, ‡ in., sterile flowers in terminal racemes or spikes, fertile flowers axillary at hase of spikes; stems rough; seed ‡ in., bearing a rown of 5-6 tubercles above the middle; leaves 3-lobed. ellow, ‡ in., sterile in racemes, fertile green, axillary; ased ‡ in.	eds,in grain, wind, floods.	ow rich la grain fie wheat.	nd, Pull,	mow, hurn plants.
llow, in., sterile in racemes, fer- ile green, axillary; seed in. long, with 6 short sharp spines; leaves inely out up.	• R	lich cultiv	ated Culti	vate late, mow bbles.