

*SALIX TENERA*, Anders.

This species was described from specimens collected by Lyall at 7,000 feet altitude on the International boundary in 1860 and was not collected in Canada again until found by the writer in 1905, probably on the same mountain on which Lyall first saw it. Second summit west of Skagit River, altitude 6,500 feet. No. 73,674.

*ABRONIA UMBELLATA*, Lane.

The record made for this species on page 148 of THE OTTAWA NATURALIST should have been credited to Mr. James Fraser instead of Mr. George Fraser.

*RANUNCULUS LOBBII*, A. Gray.

Abundant in Lost Lake near Victoria, Vancouver Island, Herb. No. 77,311. May 11th, 1908, in fine fruit. (*John Macoun*). Not before recorded north of Oregon.

*CAKILE EDENTULA*, (Bigel.) Hook.

*C. Americana*, Nutt.: Macoun, Cat. Can. Pl. I: 58.

Abundant at Long Beach, 20 miles north of Ucluelet, west coast of Vancouver Island, B.C., August 7th, 1909. Herb. No. 78,288. (*John Macoun*). Not recorded in Canada west of Lake Superior nor on the Pacific Coast north of California.

*ALTHAEA OFFICINALIS*, L.

Brother Victorin of Longueuil College reports this species to be very common in pastures at Oka, Que. We have no other record of its occurrence in Canada.

*BARTONIA VIRGINICA*, (L.) B. S. P.

The only specimens we have of this beautiful species were collected by Dr. Chas. A. Hamilton in pastures on both sides of the Feauxbourg Road just south of Mahone Bay, N.S. Dr. Hamilton reports it as growing quite plentifully over 30 or 40 acres of pasture. The soil was thin and peaty but not swampy. The plants associated with it were *Pteris aquilina*, *Myrica asplenifolia*, *Gaultheria procumbens*, etc. Herb. No. 78,291.

*AMSIACKIA LYCOPSOIDES*, Lehm.

Douglas, B.C. No. 76,744. (*Wm. Spreadborough*). Common in suitable localities on Vancouver Island but not before recorded from mainland of British Columbia.

*COLEOSANTHUS OBLONGIFOLIUS*, (Nutt.)

*Brickellia oblongifolia*, Macoun, Cat. Can. Pl. I: 207.

This plant, which is hardly distinguishable from *C. linifolius*, was collected by Lyall "on mountain slopes along the southern boundary of British Columbia" but not again until it was found by the writer in the Skagit Valley, B.C., in 1908. Herb. No.