

AAAA. Elytra rounded at tip.

b. Elytra vittate or with lateral stripes.



FIG. 5.

Pubescence silvery white; thorax and elytra with three broad pubescent brown vittae. .60-.80 in. (Fig. 5.) *candida*, Fabr.

Pubescence cinereous; head and thorax with bright yellow pubescence, six black denuded spots on thorax. Elytra with broad marginal and sutural stripe bright yellow. .37-.40 in. *puncticollis*, Say.

Pubescence grayish. Elytra with broad submarginal stripe, and sometimes also the suture narrowly yellowish-scarlet. A broad stripe on each side of thorax. Surface colour piceous, punctures coarse. .36-.60 in. *lateralis*, Fabr.

Pubescence grayish. Elytra with broad submarginal stripe (extending also along sides of thorax) yellowish-scarlet, connected with which are three oblique bands, which may reach the suture or be reduced to mere short projections, surface colour piceous, punctuation finer. .36-.52. *tridentata*, Oliv.

bb. Elytra with white pubescent spots. Surface brown.

Thorax with two white stripes, besides a narrow discal white line. Elytra each with two large white spots, sides of under surface white. .40-.80 in. *creolata*, Newm.

Thorax with two white stripes, no discal line. Elytra each with a humeral and two subsutural white spots, sometimes becoming obsolete in the male. Under side altogether or with sides white. .40-.48 in. *fayi*, Bland.

bbb. Elytra each with three small denuded spots, sometimes wanting. Pubescence dense, uniformly olivaceous or yellowish-brown. .48-.76 in. *vestita*, Say.

bbbb. Elytra with a transverse undulate fascia; surface colour piceous to ferruginous, punctuation coarse and deep. .60 in. *discoidea* (♀), Fabr.

bbbbb. Elytra unicolorous, not variegated.

Thorax with a broad denuded stripe each side of the median line. Piceous to ferruginous, under side with dense grayish - white pubescence. .40 in. *discoidea* (♂), Fabr.