

mon size and shape; the apical joint longer than the others, cylindrical, tip a little pointed, light brown; labial palpi of unusual length, longer than the head, about 3 m.m.; second joint about thrice the length of the last maxillary joint, much stronger, hairy, a little incurved, suddenly thickened above just before tip, where it is brownish; last joint as long as the second, strongly clavate before the short, fine, cylindrical tip, hairy, yellowish inside, externally brown, blackish on the club. Antennæ nearly as long as head and thorax, strong, thicker to tip, fuscous, annulated with yellowish, the two basal joints and basal half of antenna below yellowish; vertex elevated, pale yellow, on top a transversal black line interrupted in the middle, where is a black spot, and a second transversal line formed by four black spots, besides some black dots near the occiput. Prothorax a little longer than broad, narrowed before and rounded; pale with whitish villosity; two fuscous longitudinal bands, approximate and reaching the front margin; on each side, more distant, a fuscous band, ending in the transversal furrow with a black spot; below along the side margin a black stripe; mesothorax pale yellow with short black bands, forming three forks, open behind, and a similar on the hind middle part; metathorax with a black cross; sides of the thorax yellow, with a black maculose longitudinal band. Abdomen of male considerably longer than the wings, slender, with whitish villosity, intermixed with brown hairs on the apical half, fuscous below, above with a broad yellow band, divided by a fuscous line on the segments 1 to 4; a yellowish median line on the three last segments, and some lateral marks; appendages very short, cylindrical, going downwards, a little divergent, yellow with a black stripe above, strongly clothed with black hairs. I can not find below them any middle part or plate. Abdomen of female as long as the wings, blackish-fuscous, with two yellow dorsal bands from the third segment to apex; genitals light brown with a row of strong black spines; below two short cylindrical appendages with long black hairs. Legs short, pale yellow, moderately sprinkled with black, with strong black spines and on the interior of the anterior femora with white spines; tibia with a fine black ring near the base; tip of tibia and all joints of tarsi black; spurs brown, strong, as long or a little longer than the two basal joints. Wings hyaline, pterostigma white after a brown spot; costals simple, a few forked before the pterostigma; veins fuscous interrupted with pale, near the submediana fumose, forming a more or less visible longitudinal dark line; near the