

sternum. Halteres pale. Legs, coxæ and trochanters light yellow; femora light brown, becoming light yellow on the apical sixth and with a conspicuous, dark-brown, subapical ring; tibiæ dark brown; tarsi broken. Wings, whitish or subhyaline; costal cell slightly more yellowish; grey clouds as follows: At origin of Rs (largest), at stigma, at tip of Sc, along cord, along outer end of cell 1st M<sub>2</sub> and in the center of most of the cells. Venation, (see pl. III.; fig. 10); Sc long, extending far beyond the origin of Rs, Sc<sub>2</sub> at the tip of Sc<sub>1</sub>, Rs almost square at its origin and spurred (in the types), cell R<sub>2</sub> almost as far proximad as cell 1st M<sub>2</sub> (as in *F. stulta* O.S.), cell 1st M<sub>2</sub> long, longer than the veins issuing from it, basal deflection of Cu<sub>1</sub> at the fork of M.

Abdomen, tergum dark brown; caudal margins of the 7th, 8th and 9th segments more yellowish; sternum dull yellow. Hypopygium (see figs. 8, 9; pl. IV.); 9th tergite short, its cephalic and caudal margin convex its caudal half provided with a number of long hairs. Pleuræ very long, cylindrical, the tips produced into a slender lobe on the ventral side; two apical appendages, which are very short and inconspicuous, being scarcely one-third as long as the plura; dorsal appendage simple, short, slender and subchitinized, not exceeding the ventral appendage; ventral appendage double, its dorsal arm being small, triangular and with the caudal or outer face bearing a chitinized tooth, its tip produced entad and cephalad into a blunt lobe; the ventral arm is produced entad into a small lobe, with the tip evenly rounded. Viewed from the side, the pleura is broad, its ventral margin rounded at the base, at the middle of its length produced into a spatulate fleshy lobe which is directed caudad. The guard of the penis is long (extending about to the extreme tip of the pleura), and slender, broad at the base, narrowed toward the tip, the end little, if any, enlarged; the apex is very slightly notched; viewed from the side, it is seen that the extreme tip is bent ventrad; viewed from above, the guard seems to be concave, its lateral margins being more strongly chitinized. The second gonapophyses are rather long, dark brown, subrounded or scarcely pointed at the apex; at their base they are about as broad as the base of the penis guard; the lateral margin of the apophyse is produced dorsad into an incurved, chitinized flap or