

II. FACE.

	Per cent.
Disfigurement from scars of cranium (more in case of females).....	8-16
Loss of nose do	10-15
Loss of incisor teeth (females only).....	5
Defective closure of one eyelid,.....	8
“ “ both.....	16
Incurable trismus	33

III. EYES.

Loss or blinding of one (according to employment)	25-33
Loss of both eyes.....	100
Defective sight one eye.....	25
Trachoma extreme	100
Chr. conjunctivitis	15
Obstruction of lacrimal duct.....	8
Nystagmus	16

Josten's table for estimating loss of vision partial in both eyes.

RULE.—Estimate loss of central vision in each eye if over 50 per cent. (amounts below 50 per cent. disregarded).

Follow column giving the vision of one eye below V, and of the other eye to the right of V, the intersection of the column will be at the percentage of disability for both eyes.

V.	0.50	0.40	0.30	0.20	0.10	0.00
0.50	0	6.5	13.5	20.0	26.5	33.5
0.40	6.5	14.5	22.0	30.0	38.0	46.0
0.30	13.5	22.0	31.5	41.0	50.0	60.0
0.20	20.0	30.0	41.0	52.0	62.5	73.5
0.10	26.5	38.0	50.0	62.5	75.0	87.0
0.00	33.5	46.0	60.0	73.5	80.0	100.0

IV. EARS.

	Per cent.
Loss of one ear (or complete deafness one ear).....	8
Moderate deafness one ear	0
“ “ both ears unless occupation needs acute hearing)	0-8
Slight deafness both ears	0
Chr. discharge from ear	15-30

V. NECK AND VOICE.

Permanent wearing of tracheotomy tube	50
Inability to read (alexia)	75
Inability to speak (aphonia)	40
Hoarseness or whispering voice	8
Dyspnea from stenosis of larynx	33-40

VI. CHEST.

Restricted movement from distortion of chest wall, extreme	60
“ “ “ “ moderate	33
“ “ “ “ slight	8
Restriction of movement from badly healed fracture of ribs, extreme	40
“ “ “ “ medium	25
“ “ “ “ slight	8
Intercostal neuralgia	25
Lung disease, severe	100
“ medium	40
“ slight	0-8
Traumatic heart disease	0-100