

June, 1873; W. M. Wilson, 1875; Miss Pelton, 1877. Wyoming Territory, Dr. Parry, 1873. Open path near "Sheltered Falls," on path to Pike's Peak, and Clear Creek Cañon, near Georgetown, Colorado, Addison Brown, Aug., 1878.

Woodsia obtusa, TORREY.

Middlesex Co., G. E. D., 1873-5. Essex Co.: Mrs. Horner, 1874; J. R. and G. E. D., July, 1874. Lenox Mass., C. E. F.

NEPHROLEPIS. SCHOTT.

A strictly Tropical genus, our species occurring in "low rich hummocks, sometimes on the cabbage palmetto." Dr. Garber, l. c.

Nephrolepis exaltata, SCHOTT.

Indian River, Florida: Dr. Palmer (ex herb. C. E. Faxon), 1874; J. Donnell Smith, Mch., 1877. Miami, Fla., Dr. Garber, July, 1877.

DICKSONIA. L'HERIT.

Our species abundant in woodlands throughout the Appalachian range.

Dicksonia punctilobula, KUNZE.

Middlesex Co., Mass., G. E. D., 1872-4.

TRICHOMANES. SMITH.

A Tropical genus, our species occurring on dripping, or in the shaded retreating crevices of overhanging rock.

Trichomanes Petersii, GRAY.

Winston Co., on sand rock, near Sipsey River, Ala., T. M. Peters, 1873-6.

Trichomanes radicans, SWARTZ.

Winston Co., Ala., T. M. Peters, 1873. Chester Co., Ky. (ex herb. D. C. Eaton), J. Hussey; Edmonson Co. (ex herbs. C. E. Faxon and A. H. Curtiss), J. Hussey, July, 1874. Rockcastle Co., Kentucky, John Williamson, Nov., 1878.

SCHIZÆA. SMITH.

A small genus, widely diffused, our solitary species strictly local, and confined to the pine barrens of New Jersey.

Schizæa pusilla, PURSH.

New Jersey: (Pine Barrens?) Prof. Asa Gray, Donor; Little Egg Harbor, July, 1865. Crowley's Brook, near Pleasant Mills, Aug., 1874. W. H. Leggett; Atsion Bottom, F. Bourguin, Sept., 1873; Miss Eliza Hosmer, Aug., 1878.