

after the notice in my Synopsis, p. 7, where a larva from Cuba was mentioned as perhaps belonging to *Olyntha*? I think the specimen, when caught, was young; it can not be decided if the colors are fully developed; it is carded and in a delicate condition. It is of the size of *E. Salvini* McLachl., but directly to be recognized as a different species by the bifid sector of the wings.

*Oligotoma Cubana*,

Female? about 7 mill. long; dry. Head light brown, sparingly clothed with pale hairs; shape of the male; antennæ brown, densely clothed with pale hairs, 18 jointed; 1st stout, longer, 2nd very short, 3rd longer, all the following shorter, equal; the last one pointed; palpi as in the male; prothorax similar to the male, light brown; thorax and abdomen blackish brown (by exsiccation); meso and metathorax nearly equal, oblong, without any trace of wings; legs dark brown, articulation and tarsi pale; last ventral segment of abdomen not divided, lightly rounded on tip; of the appendages only the left one is present, but the tip of the apical joint is broken; the basal joint is not dilated; above, between the appendages, is a small, thin, elongated lobe.

Hab. Cuba; only one specimen carded and in bad condition, No. 136 of Gundlach's catalogue was received in 1866. If Mr. Wood-Mason's statements are to be accepted, it can be the female to the male described before. Should it be a female larva, which I scarcely believe it to be, it must belong to another and much larger species. The small larva quoted in my synopsis, p. 7, 4 mill. long, from the Museum in Berlin, was also received by Gundlach from Cuba. Probably it belongs to the same species, and is a larva.

2. *Oligotoma Hubbardi*, n. spec.

Length of the body 4 mill.; length to tip of wings 6 mill.; exp. of wings 8 mill. Male dry: Head very light brown, shining; on the sides very sparingly clothed with pale hairs; convex above, slightly narrower behind the eyes, which are black and comparatively larger than in *O. Cubana*, and less distant one from the other above; the part of the head behind the eyes a little broader than long, rounded behind, cut straight before the prothorax; antennæ (5 basal joints present) pale brown, densely clothed with pale long hairs; 1st joint short, cylindrical, scarcely longer than broad; 2nd, very small, thinner, annular; 3rd as long as the two