

The fringes are gray and flecked with brown. The hind wings and abdomen above are silvery gray, and the under side of the hind wings is the same as above, but reticulated with darker gray on the costa and the outer border. The under side of the fore wings is somewhat darker than the hind wings and shows the markings of the upper side to some extent. Described from one specimen taken in Hamilton, Ontario, and one taken at light in Amherst, Mass., Sept. 18, 1887.

I take great pleasure in naming this beautiful insect for Miss Rose Linda Davis, who has ably assisted me in my entomological work for the last three years.

NEW FORMS OF ICHTHYURA.

BY HARRISON G. DYAR.

Ichthyura multnoma N. SP.

Ground colour of primaries dark "ecru drab" (Ridg. iii., 21, * but much darker), heavily irrorate with "clove brown" (Ridg. iii., 2.) scales which almost entirely obscure the ground colour at the basal portion of the wings as well as outside the third and fourth lines. First (basal) transverse line faintly yellowish, inwardly arcuate from internal margin to median vein, then rectangularly bent towards the base and following the median vein for about 1.5 mm., when it is again bent at right angles and proceeds straight to costa. Second line before middle of wing, consisting of a line of the ground colour bounded outwardly by a narrow clove brown line, straight across wing from internal margin to costa. Third line slightly yellowish, outwardly arcuate from internal margin to vein 2, just beyond its origin, faint across cell, consisting there of a line of the ground colour apparently running straight to costa. Fourth line starting on internal margin, near origin of third line, and running slightly obliquely, but nearly straight to the costa subapically. It is undulated across the subcostal nervules, but is not broadened at costa nor marked with white, being uniformly yellowish. Outside this line subapically, an irregularly triangular rusty brown patch, and another fainter patch on the middle of external margin. A submarginal row of clove brown spots, distinct only below vein 2, being elsewhere lost in the general dark shading of the terminal half of the wing.

Thorax dark gray; the vertex of head and central thoracic patch, deep blackish brown. Abdomen, secondaries and whole under surface,

*Ridgway's nomenclature of colours, plate iii., fig. 21,