

15. *Chilosia plumosa*, Coquillett.

A single specimen taken by the writer at Glacier, Aug. 20, 1901, seems, after comparison with the type, to belong here without question. The species was described from Ormsby Co., Nevada, and has not to my knowledge been recorded elsewhere.

16. *Chilosia cyanea*, Hunter.

Port Renfrew, July 5, 1901, one specimen taken by the writer.

17. *Chilosia* sp.

Two female specimens taken at Hope Mts., July 19, 1906, and at Similkameen, July 20, 1906, by R. V. Harvey. These may be the undescribed female of some species of *Chilosia* already known from the male, but I am not able to fit them in anywhere. I hesitate to give them a new name in such a difficult group. The antennæ are plumose and dark in colour; scutellum with bristly hairs; tubercle much as in *C. tristis*, body metallic-black, with very short yellowish pile; legs black, yellowish at the knees; length, 7 to 8 mm.

18. *Chilosia* sp.

One female taken at Field, July 18, 1902, by the writer. The antennæ are lacking, and I cannot place it in any species known to me.

19. *Melanostoma coerulescens*, Williston.

Kaslo, July 2, 1903, R. P. Currie. (Determined by Coquillett.)

20. *Melanostoma trichopus*, Thompson.

Nicolum River at Hope, July 14, 1906, and Vancouver, July 7, 1906. Two specimens, by R. S. Sherman.

*(Melanostoma concinnum*, Snow.

Banff, Alberta, July 17, 1901. One specimen by R. C. Osburn.)

21. *Syrphus glacialis*, Johnson.

Vancouver, March, 9; Mission, April 4, and Hope Mts., July 19, 1906, by R. V. Harvey. The species was described from Alaska.

22. *Syrphus geniculatus*, Macquart.

Grouse Mt., July 3, 1904, and Vancouver, June 16, 1906, R. V. Harvey.

23. *Syrphus gennalis*, Williston.

Glacier, Aug. 20, 1902, R. C. Osburn, and Kaslo, June 4, 1904, J. W. Cockle.