

claws brown, as long as the spurs. Wings narrow, with fringes on hind margin, which is very little emarginate before tip; hyaline; venation pale interrupted with brown; base of numerous forks and around many transversals brown; front wings with many large rusty-brown round spots along the mediana and submediana; apical half of costal space with two series of areoles; hind wings similarly spotted, but without the large rusty-brown spots.

Length of body, male, 38 to 45 m.m.; female, 28 to 31 m.m.; exp. al., 40 to 50 m.m.

Habit., Millin, Scriven Co., Georgia; July, Morrison. I have before me two males and three females; it is a very pretty species and doubtless Rambur's *M. nebulosus*.

I had accepted this species to be *M. contaminatus* Burm., which is, with a short diagnose, only mentioned by his *M. irroratus*. Mr. Taschenberg describes a female, which is not labeled, out of Burmeister's collection. This female belongs doubtless to *M. conspersa* Rbr.; he calls the spurs of the anterior legs shorter than the first joint of tarsi, but these are probably spines. If this is really the type of Burmeister, and I have no doubt of it, the name has to be dropped, as it belongs to *M. conspersa*.

3. *Brachynemurus versutus* Walk.

Myrmeleon versutus Walk., 331, 51; Hag. Syn. Neur. N. Amer., p. 238, No. 8.

Black, very slender, faintly villous; head narrow; front shining, yellow, above a broad black band, notched in middle below, narrowly yellow around the eyes and around the base of antennæ; a black longitudinal line from the middle of the notch to the epistom, where it is enlarged and united with a larger transversal black band (which is sometimes divided into four spots) on the epistom; labrum black shining with yellow side-margins; vertex elevated, rounded, black, with a transversal yellow band, interrupted in middle, and a posterior one, representing a larger yellow spot on each side, and a smaller middle one, a little before them; antennæ of male much longer than head and thorax, about 10 m.m.; of female shorter, about as long as head and thorax, 7 m.m.; long, filiform, a little thicker to the tip, bluntly pointed; those of female a little more enlarged on tip; blackish-brown, the two basal joints shining black; palpi yellow, apical joint of the maxillary black shining, and the two foregoing black.