

Viviparus—Continued.**—plicapressus White. (1880.)**

Twelfth Ann. Rep. U. S. Geol. and Geogr. Sur. Terr., pt. 1, p. 98, pl. 28, figs. 3a, b. Wash., 1880. C. A. White.

Formation: Cretaceous.

Location: Black Buttes station, Wyo.

—plicapressus White. (1883.)

Thirteenth Ann. Rep. U. S. Geol. Sur., p. 467, pl. 24, figs. 25, 26. Wash., 1883. C. A. White.

Formation: Cretaceous.

—prudentia n. s. White. (1878)

Bull. U. S. Geol. and Geogr. Sur. Terr., vol. 4, No. 3, pp. 716, 717. Wash., 1878. C. A. White.

Formation: Cretaceous.

Location: Crow creek, Colo., ten miles above its confluence with South Platte river, northern Colorado.

—prudentia White. (1879.)

Eleventh Ann. Rep. U. S. Geol. and Geogr. Sur. Terr., p. 172. Wash., 1879. C. A. White.

Formation: Cretaceous.

Location: Crow Creek valley, Colo.

—prudentius White. (1880.)

Twelfth Ann. Rep. U. S. Geol. Sur., pt. 1, pp. 98, 99, pl. 23, figs. 5ab. Wash., 1880. C. A. White.

Formation: Cretaceous.

Location: Valley of Crow creek, ten miles above the confluence of that creek with Platte river, northern Colorado.

—prudentius White. (1883.)

Thirteenth Ann. Rep. U. S. Geol. Sur., p. 467, pl. 25, figs. 17-18. Wash., 1883. C. A. White.

Formation: Cretaceous.

—prudentius (White) Whiteaves. (1885.)

Geol. and Nat. Hist. Sur. Can., Cont. Can. Paleont., vol. 1, pt. 1, p. 24. Montreal, 1885. J. F. Whiteaves.

Formation: Cretaceous.

Location: Gooseberry canyon, St. Mary river, Pincher creek, Canada.

—raynoldsianus (M. & H.) Meek. (1876.)

Rep. U. S. Geol. Sur. Terr., vol. 9, pp. 584, 585, pl. 44, figs. 7a, b. Wash., 1876. F. B. Meek.

Formation: [Cretaceous.]

Location: Clear fork of Powder river, Mont., lower fork of same stream.

—retusus (M. & H.) Meek. (1876.)

Rep. U. S. Geol. Sur. Terr., vol. 9, pp. 578, 579, pl. 44, figs. 5a-f. Wash., 1876. F. B. Meek.

Formation: [Cretaceous].

Location: Three miles below Fort Clarke, [N.] Dak.

—retusus (M. & H.) White. (1880.)

Twelfth Ann. Rep. U. S. Geol. and Geogr. Sur. Terr., pt. 1, p. 100. Wash., 1880. C. A. White.

Formation: Cretaceous.

Location: Upper Missouri river region.

Viviparus—Continued.**—retusus (M. & H.) White. (1883.)**

Thirteenth Ann. Rep. U. S. Geol. Sur., p. 467, pl. 24, figs. 1-3. Wash., 1883. C. A. White.

Formation: Cretaceous.

Location: [Three miles below Fort Clarke, [N.] Dak.]

—reynoldsianus (M. & H.) White. (1880.)

Twelfth Ann. Rep. U. S. Geol. and Geogr. Sur. Terr., pt. 1, p. 100. Wash., 1880. C. A. White.

Formation: Cretaceous.

Location: Upper Missouri river region.

—reynoldsianus White. (1883.)

Thirteenth Ann. Rep. U. S. Geol. Sur., p. 467, pl. 25, figs. 22, 23. Wash., 1883. C. A. White.

Formation: Cretaceous.

—trochiformis (M. & H.) Meek. (1876.)

Rep. U. S. Geol. Sur. Terr., vol. 9, pp. 580-582, pl. 44, figs. 2a-c. Wash., 1876. F. B. Meek.

Formation: [Cretaceous].

Location: Ten miles below Fort Union, Mont.; also one hundred and fifty miles below Fort Union.

—trochiformis (M. & H.) White. (1880.)

Twelfth Ann. Rep. U. S. Geol. and Geogr. Sur. Terr., pt. 1, p. 100. Wash., 1880. C. A. White.

Formation: Cretaceous.

Location: Upper Missouri river region.

—trochiformis (M. & H.) White. (1883.)

Thirteenth Ann. Rep. U. S. Geol. Sur., p. 467, pl. 24, figs. 10-16. Wash., 1883. C. A. White.

Formation: Cretaceous.

—trochiformis (M. & H.) Whiteaves. (1885.)

Geol. and Nat. Hist. Sur. Can., Cont. Can. Paleont., vol. 1, pt. 1, p. 28. Montreal, 1885. J. F. Whiteaves.

Formation: Cretaceous.

Location: Souris river, four miles east of Roche Perce and Great Valley, about one hundred miles west of Wood End depot, on the forty-ninth parallel, Canada.

Volisella (Scopoli) M. & H. (1865.)

Paleont. Up. Missouri, Smithsonian. Cont. Knowl., vol. 14, No. 172, pp. 85, 86. Wash., 1865. Meek & Hayden.

—(Scopoli) Meek. (1876.)

Rep. U. S. Geol. Sur. Terr., vol. 9, p. 70. Wash., 1876. F. B. Meek.

—attenuata (M. & H.) Meek. (1876.)

Rep. U. S. Geol. Sur. Terr., vol. 9, p. 74, pl. 28, figs. 8a, b. Wash., 1876. F. B. Meek.

Formation: Cretaceous.

Location: Moreau river.