

lat. $66^{\circ} 41' N.$, long. $6^{\circ} 59' E.$, 350 fathoms, bottom temperature 0.9° coarse clay; between Norway and Iceland, Station 3, lat. $64^{\circ} 36' N.$, long. $10^{\circ} 21' 5'' W.$, in 299 fathoms, clay and sand, and Station 1, lat. $62^{\circ} 44' 5'' N.$, long. $1^{\circ} 48' E.$, in 412 fathoms, clayey bottom. Between Iceland and Faeroes: *Ingolf* Station 4, lat. $64^{\circ} 07' N.$, long. $11^{\circ} 1' W.$, 237 fathoms, temperature 2.5° ; Station 2, lat. $63^{\circ} 04' N.$, long. $9^{\circ} 22' W.$, 107 fathoms, temperature 4.8° ; west of the Faeroes, Station 44, lat. $61^{\circ} 42' N.$, long. $9^{\circ} 36' W.$, 545 fathoms, temperature 4.8° ; southwest of the Faeroes, lat. $60^{\circ} 31' N.$, long. $9^{\circ} 18' W.$, 229 fathoms; southeast of the Faeroes, lat. $60^{\circ} 10' N.$, long. $6^{\circ} 25' W.$, 620 fathoms, temperature 0.53° .

Suborder EPICARIDEA.

Family BOPYRIDAE.

Genus *Bopyroides* Stimpson.

Bopyroides hippolytes (Krøyer).

1838. *Bopyrus hippolytes* KRØYER, Kongelige Danske Videnskabenes Selskabs naturvidenskabelige og mathematiske Afhandlinger, VII, p. 306, (78), pl. iv, fig. 22.
 1840. *Bopyrus hippolytes* EDWARDS, Hist. Nat. des Crust., III, p. 283.
 1842. *Bopyrus hippolytes* KRØYER, Kongelige Danske Videnskabenes Selskabs naturvidenskabelige og mathematiske Afhandlinger, IX, p. 262.
 1849. *Bopyrus hippolytes* KRØYER, Voy. en Scand., Crust., pl. xxviii, fig. 2.
 1863. *Bopyrus hippolytes* STIMPSON, Proc. Acad. Nat. Sci., Phila., p. 140.
 1864. *Bopyroides acutimarginatus* STIMPSON, Proc. Acad. Nat. Sci., Phila., p. 156.
 1868. *Cyge hippolytes* BATE AND WESTWOOD, Brit. Sess. Crust., II, p. 230.
 1877. *Cyge hippolytes* MIERS, Ann. Mag. Nat. Hist. (4), XIX, p. 64 (14).
 1879. *Cyge hippolytes* SMITH, in Harger, Proc. U. S. Nat. Mus., II, p. 157.
 1880. *Cyge hippolytes* HARGER, U. S. Fish Comm., Pt. 6, p. 311.
 1887-88. *Cyge hippolytes* HANSEN, Videnskabelige Meddelelser fra den naturhistoriske Forening i Kjøbenhavn, p. 197.
 1895. *Cyge hippolytes* AXEL OHLIN, Akademisk Afhandling, XXII, p. 19.
 1900. *Bopyroides hippolytes* G. O. SÆRS, Crust. of Norway, II, pp. 199-200, pl. LXXXIV, fig. 2.
 1900. *Bopyroides hippolytes* BONNIER, Travaux de la station Zool. de Wimereux, VIII, pp. 373-375.
 1900. *Bopyroides sarsi* BONNIER, Travaux de la station Zool. de Wimereux, pp. 376-377.
 1900. *Bopyroides* sp. BONNIER, Travaux de la Station Zool. de Wimereux, p. 378.
 1901. *Bopyroides hippolytes* RICHARDSON, Proc. U. S. Nat. Mus., XXIII, p. 578.
 1905. *Bopyroides hippolytes* RICHARDSON, Bull. U. S. Fish Comm., XXIV, pp. 218-219.
 1905. *Bopyroides hippolytes* RICHARDSON, Bull. 51, U. S. Nat. Mus., pp. 567-568, fig. 628.
 1916. *Bopyroides hippolytes* HANSEN, H. J., Danish Ingolf Expedition, Crust. Malacost. 3, pt. 5, p. 203.

Localities recorded: This species is known to infest the branchial cavities of eight species of *Spirontocaris*, three species of *Pandalus* and one species of *Pandalopsis*.

North America:

I have examined the following specimens: Grinnell land, lat. $81^{\circ} 41' N.$, $51\frac{1}{2}$ -25 fathoms; on *S. polaris*; Cat. No. 31618, U. S. N. M., 1 specimen on *S. liljeborgii*, Georges bank, collected by the Schooner *Sultana*; Cat. No. 31619, U. S. N. M., 1 specimen on *S. polaris* from Grand Manan, New Brunswick, 3-20 fathoms; Cat. No. 39425, U. S. N. M., 1 specimen also from