

*C. sororculella*, n. sp.

Generally similar to *hjerkindella*, Thunb., but without any traces of the yellow dashes at the base of fore wings. The pale gray space in the middle of the wing is sharply limited without and within by a paler line; in the centre of this space is a large group of black and metallic scales; beyond the pale line are no black scales, but a regular, distinct, subterminal metallic line; a subcostal metallic streak in basal space. Hind wings with a white dash as in *onustana*. Two examples. Placer Co., California. June (A. Koebele); U. S. Nat. Mus., type No. 4426.

*C. occidentella*, n. sp.

Grayish brown; basal half of wing of this colour, with a curved white line across its centre. Terminal half of wing filled, except somewhat narrowly along outer margin, by a large whitish patch, irrorate with brown scales, containing above a small black patch and below a large quadrate one, cut by a whitish line transversely; silvery scales along costa basally, middle of wing, subterminally and in the black patch. Hind wings brown, immaculate. Below a faint, irregular, whitish, submarginal line on hind wings and two costal dots on fore wings. Expanse 14 mm. One male. California (coll. Beutenmüller). U. S. Nat. Mus., type No. 4428.

*C. extrinsicella*, n. sp.

Light brown, head and palpi whitish. Fore wing with basal half brown, crossed by a broad, straight, white line with a few silvery scales outwardly on costa. Terminal half of the wing nearly white except narrowly along outer margin, streaked above with longitudinal, somewhat cuneate, lines of black scales, below containing a single elongate, rounded, black patch with two groups of silvery scales; similar scales on outer border of white patch; outer half of fringe white. Hind wings pale brown. Expanse 12 mm. One male. Wisconsin. U. S. Nat. Mus., type No. 4427.

## Genus THELETHIA, Dyar.

Dyar, CAN. ENT., XXV., 301, 1893; *Thia*, Hy. Edw. (preoc. Col. 1840); Hy. Edw., Ent. Am., III., 181, 1888; Kirby, Cat. Lep. Het., I., 901, 1892.

*T. extranea*, Hy. Edw., Ent. Amer., III., 181, 1888; Smith, List. Lep. Bor. Am., No. 956, 1891; Dyar, CAN. ENT., XXV., 301, 1893.

I include this genus here tentatively. I have not seen a specimen in six years, and my old notes give only the venation. It may be a *Tineid* near *Incurvaria*.