

its post. extremity, and 1 inch at its ant. extremity, and of a pear shape. The skin surrounding the ulcer, and for about 3 inches above it, was highly inflamed and exceedingly tender to the touch, with slight effusion beneath. He appeared perfectly healthy in every other respect, but complained of anorexia. Linseed poultices were applied upon first entrance.

27th Sept.—Was ordered red wash to be constantly applied to the part:—R Liq. Arseni, chl. 3 vi., Aqua 3 iv. Dose, 3 ii. in water, three times each day. This treatment continued until Oct. 13. He stated then that his appetite was much better, and altogether he felt much improved. The ulcer was healthy-looking, all the inflammation had subsided, and new skin to the extent of  $\frac{1}{4}$ th of an inch had formed all around the sore. But it was decided in consultation that the stump could never be useful to him, and he accordingly submitted to a re-amputation, which was performed by Dr. Wright in the circular manner, just below the tuberosity of the tibia, Oct. 14th.

*State of Case after Amputation.*—Oct. 15. Patient very restless and excited; an an-dyne draught was administered during the night. Ordered Imperial as a drink. Acet. Opii H. S.

Oct. 16. Patient's face is flushed, and he is still restless: progress is satisfactory.

Oct. 19. Pulse small and quick; wound doing well, small collection of pus in the depending part, is very tender, but not inflamed, (straps and bandage). Ordered 5 oz. of wine, and ditto. of brandy; R Quinae Sulph. xv. grs., Acid Sulph. Arom. 3 ii., Ether Sulph. Spts. co. 3 ii., Aqua 3 vi. 3 ss. three times each day.

Oct. 20. Omit Imperial on account of some intestinal distress. Ol. Ricini 3 ii. Wound looks well; most of it has healed by first intention. No pain of consequence, and little irritability.

Oct. 23. Was very excitable and restless; rolled much in bed, and between the hours of 7 and 8 in the evening struck the stump against the cradle, and hemorrhage set in. Blood came away in a small jet, a little more than the size of a thread, reaching the height of about 2½ feet about 8 oz. of blood was lost. Dr. Craik, House Surgeon, was sent for; he applied the tourniquet to the femoral, and the bleeding ceased. Dr. Wright arrived: took off the bandages, and reapplied them. Bleeding did not recur, and no untoward event has been felt since.

Oct. 25. Wound looks healthy, and patient better and stronger; ordered mutton chop and potatoes.

Oct. 28. Ordered clothes.

Nov. 1. All the ligatures have safely come away. The line of the