Black Pigeon Hawk—Falco columbarius sucklei.
One record, February 7, 1914.

Desert Sparrow Hawk—Falco sparverius phalana.

One winter record, January 22, 1917.

LONG-EARED OWL-Asio wilsonianus.

Common resident.

SHORT-EARED OWL-Asio flammeus.

Common winter resident; a few remain to breed.

GREAT GREY OWL—Scoptiaptex nebulosa nebulosa.
Scarce winter resident.

RICHARDSON'S OWL—Cryptoglaux funerea richardsoni Scarce winter resident.

SAW-WHET OWL-Cryptoglaux acadica acadica.

Resident; not common.

MACFARLANE'S SCREECH OWL—Otus asio macfarlanei.

Common resident.

GREAT HORNED OWL-Bubo virginianus.

The breeding form is consistently dark with the heavily mottled feet of "saluratus." In the winter a much lighter, more ruescent form occurs. I have never taken anything approaching subarcticus. Snowy Own—Nyctia nyctia.

Winter resident; not common. During the winter of 1916-17, when the big migration of this species occurred on the Pacific coast, they were slightly more common here.

HAWK OWL—Surnia ulula caparoch.

Winter resident; not common. Breeds in the high mountains in southern Okanagan.

Pygmy Owl.—Glaucidium gnoma gnoma.

Common resident.

Belted Kincfisher—Ceryle alcyon.

A few generally winter.

ROCKY MOUNTAIN HAIRY WOODPECKER—Dryobates villosus monticola. Common resident.

Batchelder's Woodpecker—Dryobates pubescens homorus.

Common resident.

WHITE-HEADED WOCDPECKER—Xenopicus albolarvatus.
Two records, December 20, 1911; January 24, 1914.

ARCTIC THREE-TOED WOODPECKER—Picoides arcticus.

Resident in the Canadian zones.

ALASKA THREE-TOFO WOODPECKER—Picoides americanus fasciatus. Resident in the Canadian zone.

NORTHERN PILEATED WOODPECKER—Phloeotomus pileatus abieticola. Common resident.

RED-SHAFTED FLICKS-Colaptes cafer collaris.

A few always winter.