

abundant; head broad; clypeus not much produced, the lower half black; supraclypeal area brassy; front and sides of face well punctured, the middle of front as closely punctured as possible, not striate; antennæ black; cheeks shining; mesothorax shining, strongly and rather closely punctured all over; scutellum closely, more finely punctured; area of metathorax strongly sculptured all over with anastomosing ridges, producing a finely reticulate effect; tegulæ rufous with paler margins, impunctate; wings hyaline, not reddish; stigma and nervures reddish testaceous; hind spur with three long blunt spines, the basal one remarkably long; first abdominal segment smooth and shining, though minutely punctured; the other segments duller and more punctured, those beyond the second pruinose with pale hair.

South Island, Coronado Islands, L. Calif., Aug. 21 (*T. and W. Ckll.*). Five females, visiting *Compositæ*. One would not take this for a southern or desert species; it rather has the aspect of those of the mountains and of the north. It falls near to *H. perpunctatus* Ellis, but has the mesothorax more coarsely and less densely punctured, wings not reddish, area of metathorax shorter, etc. A related but quite distinct form, which Mrs. M. D. Ellis has named in manuscript as a subspecies of *H. perpunctatus*, occurs in Southern California. I have named this species and the next after the naturalists to whom we are indebted for an excellent account of the birds of the Coronado Islands.

***Halictus daggetti*, n. sp.**

Male—Length about 5.5 mm.; head and thorax olive green, abdomen black; pubescence white; clypeus moderately produced; face and front hairy; antennæ very long (reaching middle of metathorax), flagellum rather thick, ferruginous beneath, narrowly so on the apical part; inner orbits submarginate; mesothorax and scutellum brilliantly shining, well punctured, sparsely on disc of mesothorax; area of metathorax sculptured essentially as in *H. grinnelli*; tegulæ shining piceous; wings milky-hyaline, stigma testaceous bordered with fuscous, nervures fuscous, second s. m. narrower above; mesopleura closely punctured; legs black, with pale hair; abdomen shining, but finely and rather closely punctured, with erect pale hair, especially conspicuous at sides; the suture