

ISOPODA.

Lygia dilatata Stimpson.

One specimen from near Victoria, V.I.

Synidotea nodulosa Harger.

Idothea nodulosa Krøyer, Naturhist. Tidssk., II., ii., p. 100, 1846; in Gaimard, Voyage en Scandinavie, pl. 26, fig. 2, 1849.

Synidotea nodulosa Harger, Amer. Jour. Sci., III., xv., p. 374, 1878; Proceedings United States National Museum, 1879, p. 160, 1879.

Two specimens from 111 fath., Dixon Entrance, Q.C.I. It has been found in the Atlantic from George's Banks and Nova Scotia to Greenland, but has not been recorded heretofore from the Pacific. The specimens were determined by Mr. Harger.

Spharoma sp.

A small species from Dolomite Narrows, Q.C.I. It is apparently quite distinct from *S. Oregonensis* Dana and from *S. amplicauda* Stimpson, the only species, as far as I know, described from the north-west coast of America.

Tanaïs? sp.

There are two dry specimens of a small Tanaid from 15 to 18 fath., Virago Sound, Q.C.I.

CIRRIPEDIA.

Tetraclita porosa Darwin.

Near Victoria, V.I.

Lepas anatifera Linné.

Near Victoria, V.I.