

General colour a very bright grass green. Fastigium tipped with dull yellow, which extends half way down the sides. Labrum and apical segments of all the palpi a rose red tinged with violet. Tarsi somewhat infuscated. Antennæ and apical third of ovipositor reddish-brown.

Measurements.—Female—Length of body, 27 mm.; of fastigium in front of eye, 2.75 mm.; of pronotum, 7 mm.; of tegmina, 37 mm.; of hind femora, 20 mm.; of hind tibiæ, 19.5 mm.; of ovipositor, 19 mm.

This handsome species of *Conocephalus* belongs to the same group as *C. robustus* and *C. crepitans*, Scudder, but is smaller and of a more uniform and brighter green than either of those species, besides having shorter legs, ovipositor, etc. It is described from a single female taken October 24, from the fallen grasses on the margins of a large low-land pond in Vigo county. This pond is surrounded on all sides by heavy timber, and its margins have yielded a number of interesting Orthoptera found nowhere else in the county. Among them are *Leptysma marginicollis*, Serv., *Paroxya atlantica*, Scudder, *Anaxiphus pulicarius*, Sauss., *Phylloscirtes pulchellus*, Uhler, and *Xiphidium nigro-pleurum*, Bruner. The first four mentioned are insects of a southern range, and perhaps *C. palustris* will in time be found to be more common southward.

ORCHELIMUM, Serville.

Orchelimum indianense, nov. sp.

A slender-bodied insect, with a dark median streak down the face, and having the posterior femora unarmed beneath. The cone of the vertex is short, rather narrow, with a rounded apex. The tegmina narrow, tapering, a little shorter than the wings, and of a delicate, almost gauze-like, texture. Posterior femora slender, shorter than the closed tegmina. Anal cerci of male of medium size, longer than the subgenital plate, tapering to a dull point; the basal tooth short, with a broad base and a very sharp point. The ovipositor of female of less than average width and length, the apical half with a gentle upward curve.

Colour of dried specimens: Tegmina and wings a transparent whitish, tinged with green on the front or lower, longitudinal nerves; the cross nervules of the latter darker. Sides of pronotum and abdomen, and all the femora, light green; the tibiæ and tarsi of a brownish hue. Face yellowish-white, with a dark fuscous stripe, the width of the labrum, starting with the mouth and passing upward to the vertex, where it narrows to the width of that organ; then, broadening on the occiput, it passes back to the front border of the pronotum, where it divides into