

are lost and the wing is longitudinally shaded with whitish on median space along internal margin, and diffusely beyond the reniform. New York.

*Agrotis Hospitalis*, n. s.

Allied to *baja* by the black mark, inaugurating s. t. line. Yellow-brown. Half-line black, single; t. a. line black, single, broken, irregularly trembled or denticulate; a dot between these lines on median vein. T. a. line marked on costa: t. p. line appearing as a pale even shade, preceded by broken black scallops not evident, the line itself. Orbicular large, round, paler than the wing, ringed with black, the costa above it pale like the spot itself. Median field shaded with darker brown behind. Reniform ringed with black, kidney-shape, paler than wing, dark or black inferiorly. An interrupted dark terminal line. Fringe concolorous. Hind wings rather pale, silky, concolorous. Palpi dark brown at the sides, tipped with pale brown like front in color. Mr. Hill's collection, N. Y., July at sugar.

*Gortyna Impecuniosa* Grote.

This is a dark species, almost recalling *nebris*. Dark purple brown with the median field rusty. Stigmata darker-ringed, concolorous, all indicated. A faint apical yellowish patch, inaugurating s. t. line, which is lost or fragmentary. Hind wings dark fuscous. A sharp tuft behind the collar. Mr. Hill's collection, Centre, Sept. 25.

This is allied to *Harrisii*, etc., but, I think, distinct. It is the most obscurely marked of any of the purple-brown species. One male. Size of *Harrisii*.

*Rheumaptera Immediata*, n. s.

Pale gray with all the transverse markings indistinct except the two black shaded median lines distinct, enclosing the reddish brown median space. Inner line *curved*. The lines are rather neat. The outer with a strong submedian curve, projected somewhat narrowly at median vein. Costal dots distinct. Basal line fine, not very noticeable. Beneath with an evident angulate common outer line, sub-punctate, terminal space shaded outwardly and especially over apices with brown. Discal dots and, on fore wings, a costal dot, marking inner line. Hind wings above pale, with a mesial sinuous line. Under surface darker than upper. Expanse 26 mil. Two specimens, Mr. Hill, New York.