

A FEW NEW SPIDERS.

BY NATHAN BANKS, SEA CLIFF, N. Y.

Micaria gentilis, n. sp.

Length, 3.6 mm.; ceph.: long, 1.4 mm.; broad, 1 mm. Cephalothorax, sternum, femora i. and ii., blackish; rest of legs i. and ii. pale yellowish; legs iii. and iv. yellow-brown, the femora scarcely darker; abdomen black above, with a narrow white band across the middle and a white spot on each anterior lower side; venter pale. Cephalothorax broad, in ♂ a little narrower; posterior row of eyes procurved, the P. M. E. oval, fully their diameter apart, and about as far from the P. S. E.; anterior row strongly procurved, the A. M. E. fully their diameter apart, and about as far from the larger A. S. E. Sternum oval, pointed behind; legs of moderate length, femora i. and ii. stouter than others, femora iii. slightly excised before the tip behind. Abdomen not constricted, quite broad, somewhat depressed, epigynum appearing much like *M. montana*, Em., but the openings are farther apart and more oblique. Tibia of ♂ palpus has short projection at tip on the outer side: the bulb is triangular in side view, the red parallel marks are along the outer edge; near the middle is a short tube.

Several specimens from Franconia, N. H. [Mrs. Annie T. Slosson]. Related to *M. perfecta* from Colorado, but larger and with a broader sternum.

Scius montanus, n. sp.

Length, 2.1 mm.; ceph.: long, 1 mm.; broad, .6 mm. Jet black, shining, almost coppery; extreme tips of palpi, tips of maxillæ, a spot on each coxa and trochanter, and an elongate spot on the femur, pale; the tarsi infuscated; pale dots on legs at origin of hairs. Cephalothorax long, moderately low, nearly flat, sides almost parallel. Eye region one-third broader than long, occupying not much over one-third of the cephalothorax; a trifle broader in front than behind; eyes of second row full as close to the dorsal eyes as to lateral eyes; A. M. E. large, distinctly separated, plainly farther from the S. E. Sternum one-fourth longer than broad, broadest in middle, pointed behind, truncate in front; coxæ i. separated by full width of lip; legs short, fourth pair longest, femora i. thicker than others; only a few indistinct spines, those on metatarsi iv. are at apex. Abdomen barely wider than cephalothorax and but little longer, pointed behind. Body and legs clothed with scattered black hairs. The ♂ palpi short, the tibia with a short process on the