

6. It is necessary in all cases to begin with a weak current, slowly increased in intensity, and to stop short of producing more than slight pain, to gradually accustom the patient to its effects and so overcome all physical and moral resistance.

7. Gradually in two or three sittings the intensity of the current is to be increased until in the majority of cases 100, 150 and if necessary 200 milliampères have been attained. The intensity is to be proportioned and regulated by the tolerance of the patient and the extent, gravity and duration of the lesion.

8. The duration of the application, from five to ten minutes, must, like the intensity, be regulated by the extent of effects it is desirable to obtain.

9. The sittings may be held weekly or tri-weekly, according to necessity, and the physician must regulate the number and intervals of the applications according to the urgency and necessity for intervention.

10. The patient must be made to rest in all cases several hours after the application. Such rest is necessary for the safety and efficacy of the method.

11. Antiseptic vaginal injections of corrosive sublimate or carbolic acid are to be prescribed for administration morning and evening.

This simple and safe treatment, a true therapeutic hysterometry (veritable hysterometrie thérapeutique), is none other than a molecular galvano-chemical curetting, acid or basic, according to the case, which excites the formation of a new mucosa and constitutes a kind of intra-uterine issue, the action of which we may prolong or vary as we please. Its beneficent effects which I have verified in a large number of patients are speedily obtained from the first sittings, become rapidly more marked, and soon lead to a cure. Confinement to bed or any additional treatment is unnecessary. As compared with surgical curetting, it has the advantage that it can be localized and regulated as regards dose or amount, at pleasure, according to the necessities of the case.