

two or three days with the following : Flax seed meal, mixed with a solution of two drachms sulphate zinc to a pail of water ; bathe frequently with glycerine or a solution of zinc, or a solution of chloride of lime.

*Water Farcy.*—Give one of the following powders night and morning in the feed : Sulphate of iron two ounces, nitrate of potassa one ounce, pulverized gentian one ounce, pulverized ginger six drachms, anise-seed ground half ounce, mix and divide into eight powders.

*Founder.*—If the animal is in full condition, bleed freely from the feet and give the following : Barbadoes aloes six drachms, croton oil six drops, pulverized ginger one drachm, pulverized gentian two drachms, mix with water in form of ball ; foment the feet well with hot water and then poultice with flax seed meal for several days ; give in the water, every six hours, extract of belladonna one drachm.

*Shoulder Strain.*—Bleed freely from the plantar vein running down upon the inside of the front legs. Foment the shoulders well with hot water, if the case is a recent one ; if of long standing, a seton will be more effective. The following liniment will be a useful application ; Sweet oil one pint, spirits of hartshorn three ounces, spirits of turpentine two ounces ; mix all together ; shake well before using.

*Capped Hock.*—Blister with tincture of iodine, or iodine ointment is useful.

*Bone Spavin.*—When there is external enlargement, active blisters should be applied over the part. Liquid blister : powdered croton seed half ounce, powdered cantharides one ounce, oil of turpentine one pint, olive oil one pint, mix all together and shake well before using.

*Curbs.*—In recent cases, the part should be bathed with tincture of iodine once a day. Take a little blood from the saphena vein on the inside of the hind leg, above the hock. Should this not succeed, blisters must be resorted to.

*Blood or Bog Spavin.*—Use cold water compresses placed upon the joint, for six or eight weeks, by means of a leather socket, made to fit. Old woollen or muslin cloth is the best.

*Palpitation of the Heart.*—The worst cases yield in two hours to the following simple treatment : Divide one drachm of digitalis into five powders, and give one every fifteen minutes on the tongue.

*Inflammation of the Brain.*—Open the jugular vein as quickly as possible ; it should be done before the mad stage comes on, or it is too late to be of much service. Open the bowels freely—give the following : Barbadoes aloes one ounce, croton oil ten drops ginger one drachm, gentian one drachm ; mix with molasses or honey. Give tobacco-smoke injections, if convenient ; or, soap and water will answer the purpose. Give, on the tongue, every two hours, ten drops tincture of aconite ; until eight doses have been given, and then stop giving the aconite ; give cold water to drink and apply cold-water bandages to the head ; or, bags of ice would be better. Give no food for twelve hours after relief is obtained.

*Stomach Stagers.*—Give the purging ball recommended in inflammation of the brain and bleed freely from the jugular vein ; give no food for forty-eight hours. This is all the treatment the animal requires.