

10 CONTRIBUTIONS TO NORTH AMERICAN ICHTHYOLOGY—I.

Rafinesquian genera.	Type species.	Modern genera.
Apomotis *	cyanellus.	Apomotis.
Notemigonus	auratus.	Notemigonus (<i>Stilbe</i>).
Amphiodon	"alveoides" (<i>alosoides</i>).	Hyodon.
Ambiodon	bubalus (teeth of <i>grun-</i> <i>niens</i>).	Ichthyobus and Haplo- donotus.
Cycleptus	nigrescens.	Cycleptus.
Pylodictis	limosus.	Pelodichthys (<i>Hoplade-</i> <i>lus</i>).

1820.

Stizostedion *	salmonea.	Stizostethium
Lepibema *	chrysops.	Roccus Mitchell, 1817.
Pomacampus *	nigropunctata.	Stizostethium.
Ichthelis	auritus L.	Lepiopomus.
Telopomis *	—	Apomotis (<i>Erythrus</i> C. & V.).
Aplites *	pallida.	Micropterus Lac.
Nemocampus *	flexuolaris.	Micropterus Lac.
Dioplites *	salmonea.	Micropterus Lac.
Ambloplites *	ichtheloides.	Ambloplites.
Aplesion *	calliura.	Micropterus Lacép.
Diplesion *	blennioides.	Diplesium (<i>Hyostoma</i> Ag., 1854).
Pomolobus	chrysocloris.	Pomolobus.
Dorosoma	notata.	Dorysoma (<i>Chatoëssus</i> Cuv. & Val., 1829).
Clodulus *	clodulus (Le S.)	Hyodon Le S.
Minnilus	—	Nototropis.
Alburnus	alburnus L. (Euro- pean).	Alburnus (Heckel emend., 1843).
Phoxinus	phoxinus L. (Euro- pean).	Phoxinus (Agassiz emend., 184-).
Dobula	dobula L. (European).	? Squalius Bonaparte, 1841.
Hemiplus (181-).	— (European ?).	—.
Luxilus	chrysocephalus.	Luxilus (<i>Hypsilepis</i> Baird, 1854).

Rafinesq
Chrosom
Semotilus
Rutilus
Plargyrus
Pimepha
Moxostor

Ictiobus
Carpioide
Teretulu
Enyrtor
Decactyl
Cycleptu
Ictalurus
Elliopt
Ameiuru
Ilietis *
Leptops
Opladelu
Picorellu
Cylindro
Atractos
Sturio *
Sterletu
Dinectur
Pegedio
Proccero

II.—LII

I give
Rafines
the spe
my ide