

4. PEACH. *Brown fruit rot, leaf blight, plum curculio and peach curl.*
 Spray at first as in case of plum, but after growth begins make Bordeaux mixture only of half strength.

Fourth spraying: Ammoniacal copper carbonate, if any danger of disfiguring the fruit with Bordeaux mixture.

In spraying peach trees for curl leaf or peach rot Bordeaux mixture full strength may be used while the tree is dormant. After growth begins use it only half strength.

5. CHERRY. *Aphis, slug, brown rot and leaf blight.*

First spraying: Bordeaux mixture and Paris green as the buds are breaking; if *aphis* appears use kerosene emulsion alone or whale-oil soap.

Second: Bordeaux mixture and Paris green as soon as the blossoms fall.

Third: Bordeaux mixture and Paris green ten to fifteen days after.

6. GRAPES. *Mildew, black rot, flea beetle, and leaf-eating insects.*

First Spraying: Bordeaux mixture and Paris green when the leaves are one inch in diameter.

Second: Bordeaux mixture and Paris green when flowers have fallen.

Third and fourth: Bordeaux mixture at intervals of ten to fifteen days.

Paris Green alone when the beetle is attacking the buds in the spring, or when the larva is attacking the leaves in summer.

7. RASPBERRY. *Anthraxnose, leaf blight and saw-fly larvæ.*

First: Bordeaux mixture and Paris Green just before growth begins.

Second: Bordeaux mixture and Paris green about when first blossoms open.

Third spraying: Bordeaux mixture when the fruit is gathered.

8. CURRANT AND GOOSEBERRY. *Worms and mildew.*

First spraying: Potassium sulphide, Bordeaux mixture and Paris green as soon as the leaves expand.

Second spraying: The same ten to fifteen days later.

For worms alone, hellebore or Paris green will be effective.

9. TOMATO. *Rot and blight.*

Spray with Bordeaux mixture, as soon as rot or blight appears, for three times, if necessary, at intervals of ten to fifteen days.

10. POTATO. *Blight and beetles.*

First spraying: Paris green as soon as the beetles appear (one pound to 100 gallons of water).

Second spraying: Bordeaux mixture and Paris green when plants are six inches high.

Third and fourth sprayings: Bordeaux mixture at intervals of ten to fifteen days, if necessary.

Spraying with Bordeaux mixture will prevent the blighting of the plants and the rotting of the tubers.

11. CABBAGE WORM.

Pyrethrum applied in solution (one ounce to four gallons of water) or dusted on (one part pyrethrum to seven parts flour) for the cabbage worm.

12. STRAWBERRY. *The rust or leaf-blight.*

Bordeaux mixture, when it can be applied without disfiguring the fruit, will control this disease. Apply at intervals of two or three weeks on new beds after they begin to make runners.