

NICOTIANA, L. Tobacco.

1623. *N. rustica*, L. (Wild Tobacco.)

Introduced. A coarse weed with green flowers.

SCROPHULARIACEÆ.—Fig-wort Family.

VERBASCUM, L. Mullein.

1625. *V. Thapsus*, L. (Common Mullein. Flannel-leaf.)

Introduced. Very common. July—1. (B)

1627. *V. Blattaria*, L. (Moth Mullein.)

Introduced. Rare. Near Cummings Bridge (*R. B. Whyte*), near the St. Louis Dam. Beechwood. There are two forms of this plant—one with yellow flowers, probably identical with the English plant, and a form with larger white flowers tinged with purple. It is probable that this last is an American variety.

LINARIA, Juss. Toad-flax.

1629. *L. vulgaris*, Mill. (Butter and Eggs.)

Introduced. Common. July—1. (B) The flowers vary much in depth of colour, from almost white to orange.

————— var. *Peloria*. This is a rare monstrous state with a regular 5-cleft border to the corolla, 5 spurs and 5 stamens. Plants collected in Metcalfe street produced racemes with every flower of this nature for several years.

CHELONE, L. Turtle-head.

1637. *C. GLABRA*, L.

In bogs and wet meadows. Common. July—1. (B)

PENTSTEMON, Mitchell. Beard Tongue.

1647. *P. PUBESCENS*, Solander.

Rocky banks. Rare. Hull. Little Chaudiere. Aylmer. July—1. (B)

MIMULUS, L. Monkey-flower.

1654. *M. ringens*, L.

In ditches and low ground. July—1. (B)