

middle of the clypeus with a faint golden tinge. Wings subhyaline, tegulae and nervures testaceous. Body very finely punctured, the punctures more sparse upon scutellum; sides of thorax delicately striate longitudinally; metathorax divided by a median impressed line, delicately striate, the longitudinal striae of the base curving to meet those of the sides, the striae of the posterior face transverse. Basal segment of the abdomen more coarsely punctured than the following segments. Enclosure on sixth segment flat and broad, coarsely punctured, the margin smooth.

♂. Length 10 mm. More slender, tubercles oscillated with black, spot behind them smaller, line on collar more widely interrupted, only one dot at the extreme lateral anterior angle of scutellum, no spot on metathorax, metathorax more coarsely striate, fascia on the third segment of abdomen entire, the fascia on fifth segment narrow and a similar fascia on sixth segment. Apical segment with a longitudinal impression. Antennae twelve-jointed, the third joint excavated at the base beneath, the fourth and fifth joints slightly excavated beneath; the sixth joint arcuated, much excavated beneath, thickened at the apex, the seventh joint obliquely attached and short like the following joints. The two basal joints of anterior tarsi white, a little dilated externally, the three apical joints black; first joint of intermediate tarsi short, much thickened beneath, especially towards the apex; second joint arcuated, produced at the apex beneath.

Waterbury, Conn., ♂ Aug. 23rd; ♀ Aug. 24th.

Allied to *C. dilactus* Cress., but in that species there is no yellow spot on the metathorax, the abdominal fasciae are broader, and in the ♂ the third joint of the anterior tarsus is white.

*BLEPHARIPUS UNICUS*, n. sp.

♀. Length 5 mm. Black; tips of the mandibles, the tegulae, spurs of posterior tibiae and extreme base of the first joint of posterior tarsi, the last joint of posterior tarsi, the tips of all the coxae and trochanters and the tips of the posterior femora and tibiae, piceous. Scape beneath, dot on first joint of flagellum, the tubercles, the four anterior tibiae excepting a black spot beneath, and the tips of the four anterior femora, yellow. The four anterior tarsi, excepting the fulvous apical joint, and the base of the posterior tibiae, whitish. Clypeus black, covered with a silvery pile; flagellum fulvous beneath. Thorax beneath and the abdomen with short scattered pubescence. The abdomen excepting the rufo-piceous