

a slender point in front, behind is a transverse area pointed in front, and in front are two dark spots connected to the posterior area by a reddish line. Taken from graves, Washington, D. C. (Dr. Motter.)

Nesticus cavicola, n. sp.

Length ♂, 1.5 mm. Wholly pale whitish, clothed with long bristly hairs. Cephalothorax rather short and broad; six eyes, sub-equal in size, A. M. E. not visible; posterior row nearly straight, the P. M. E. farther apart than from the P. S. E.; S. E. touching; sternum broad, sides rounded; legs long, all femora longer than the cephalothorax, hairs on legs longer than the diameter of the joints; abdomen pointed behind, one and a half times longer than broad. Femur of male palpus rather long, palpal organ large, a projection of tibia broadest near tip and bifid, one branch is cleft; the style is long, curved around tip of bulb, from the tip of bulb there projects outward a pointed spine, and below is another projection tipped with a short black hook. From a cave, Chiricahua Mts. (Wood Canon), Ariz., June, 1897. (H. E. Hubbard.)

Erigone albescens, n. sp.

Length ♀, 1.8 mm. Cephalothorax, legs, mandibles, and sternum uniform yellowish, abdomen uniform whitish gray; eyes on black spots. Head rather elevated; posterior eye-row procurved, the P. M. E. about as far from each other as from the equal P. S. E.; A. M. E. small, and close together. Mandibles of moderate size, vertical, armed along their lower front margin with several teeth, and behind with a row of denticles, sternum broad, sides rounded, blunt pointed behind; legs of moderate length, with many hairs and a few spines, one above on tibia IV.; abdomen oviform, clothed with scattered stiff hairs, which arise from minute yellowish dots; epigynum shows two curved dark lines approaching each other from behind, and a darkish spot outwards from their tips. From the inside of coffins in graves opened during the transfer of a cemetery, Washington, D. C. (Dr. Motter.)

Philodromus pacificus, n. sp.

Length ♀, 4.1 mm. Cephalothorax yellowish, darkest on the sides, which are densely mottled and lineate with red-brown, a white \vee mark on the middle with its apex near the dorsal groove, dorsum of abdomen white, showing two basal pairs of yellowish muscular spots, sides of abdomen broadly suffused with red-brown from base to tip, venter whitish; femora, patellæ, and tibiæ pale reddish brown above, lighter below, distal joints yellowish. Sternum light yellowish, the body is full,