

III₄, III₅ shorter, subquadrate, with vein IV₃ beyond its middle ; m. p. 6, a little longer than galea ; l. p. simple, 5:1 ; glossa lance-linear, acuminate, shorter than mentum ; mandibles bidentate ; femorilegid, scopa plumose, a simple curl on base of femur ; 17 mm ; *Paranomia Nortonii* in.....(3) Nomiidæ.

Tegulæ ordinary 1.

1. Labrum free from mandibles, as large as clypeus, shorter than wide, transversely striate, without basal process ; cell III₁₊₂ pointed on costa ; 2 submarginal cells, subequal ; femorilegid, scopa simple, a thin floccus on trochanter plumose ; mandibles bidentate ; m. p. 6, longer than galea or stipes ; l. p. simple, 6:4, longer than mentum ; glossa lance-linear, acuminate, longer than mentum ; nervures pale ; thorax with dull ochraceous pubescence ; segments with apical margins pale testaceous ; 7-8 mm ; *Halictoides marginatus* in.....(4) Dufouroidæ.

Labrum ordinary 2.

2. Hind tibia and metatarsus broad, with dense simple hairs, white on former, black on latter ; pollen mixed with honey carried mainly on anterior faces of both joints ; cell III₁₊₂ pointed on costa ; 2 submarginal cells, III and III₄₊₅, subequal ; vein *m* about equals V₁ ; mandibles bidentate ; m. p. 6, shorter than galea l. p. simple, 2:1 ; abdomen closely and coarsely punctured ; 9 mm ; *Macropis steironematis* in.....(5) Macropididæ.

Hind tibia and metatarsus ordinary 3

3. Cell III₁₊₂ truncate ; 2 submarginal cells ; facial foveæ present, narrow, glabrous ; mandibles simple ; labrum with a median area ; crurilegid, pollen mixed with honey carried mainly on anterior face of tibia(6) Panurgidæ.

Cell III₁₊₂ pointed on or near costa, acuminate beyond vein III₄ ; usually 3 submarginal cells, III = III₄ + III₅, or nearly ; vein *m* shorter than V₁ ; femorilegid ; m. p. 6 ; l. p. simple, at most 1 = 2-4 ; glossa shorter than mentum ; vein *a* rarely a little before V₂ . . . 4.

4. Vein *a* strongly bent or arcuate ; *m* quite oblique to line of V₁ ; IV₃ beyond middle of cell III₅ ; no facial foveæ ; scopa plumose, femur with long hairs posteriorly* ; m. p. longer than galea, shorter than stipes ; glossa lanceolate or lance-linear, acuminate(1) Halictidæ

*Scopa wanting in inquilines.