

Erebia discoidalis Kirby.

Satyrus alope Fabr. vars. *nephele* Kirby and *olympus* Edw.

Chionobas varuna Edw.

LYCÆNIDÆ.

Thecla acadica Edw.

Thecla calanus Hbn.

Thecla liparops Bd.-Lec.

Thecla augustus Kirby.

Thecla titus Fabr.

Thecla heathii Fletcher. There were no specimens under this name in the collection. The species was described in 1903 from a single female taken by Mr. Heath at Cartwright, about twenty-five years previously. As Dr. Skinner has suggested, it may prove to be an aberration.

Chrysophanus thoe Bdv.

Chrysophanus helloides Bdv.

Lycæna sæpiolus Bdv.

Lycæna afra Edw. The form stood as *couperii* Grt., but Mr. Wallis tells me that it is now called *afra* (teste, Dr. Skinner).

Lycæna rustica Edw. One female standing as *aquilo*. On comparison with Calgary material the specimen appears to be normal female *rustica*.

Lycæna melissa Edw.

Lycæna pseudargiolus Bd.-Lec. and vars. *lucia* Kirby and *neglecta* Edw.

Lycæna amyntula Bdv. Heath appeared to have separated two series as *amyntula* and *comyntas*, but the separation did not seem justified.

PAPILIONIDÆ.

Pieris protodice Bd.-Lec.

Pieris napi Linn. var. *oleracea-æstiva* Harr.

Pieris rapæ Linn.

Nathalis iole Bdv.

Colias cæsonia Stoll.

Colias eurytheme Bdv.

Colias eriphyle Edw.